

=> d his full

(FILE 'HOME' ENTERED AT 08:25:08 ON 24 JUN 2005)

FILE 'REGISTRY' ENTERED AT 08:25:33 ON 24 JUN 2005
ACT AUD807F0/Q

L1 STR

L2 STR L1
D QUE L2
L3 SCR 1841
L4 SCR 2039 OR 2041 OR 2050 OR 2049 OR 2053 OR 2052 OR 2048 OR 204
L5 0 SEA CSS SAM L2 AND L3 NOT L4

FILE 'HCAPLUS' ENTERED AT 08:30:49 ON 24 JUN 2005

L6 1 SEA ABB=ON PLU=ON (GB99-17793# OR WO2000-GB2903#)/AP,RPN

FILE 'REGISTRY' ENTERED AT 08:32:01 ON 24 JUN 2005

FILE 'HCAPLUS' ENTERED AT 08:32:03 ON 24 JUN 2005

L7 TRA L6 1- RN : 65 TERMS

FILE 'REGISTRY' ENTERED AT 08:32:03 ON 24 JUN 2005

L8 65 SEA ABB=ON PLU=ON L7
L9 16 SEA ABB=ON PLU=ON L8 AND NR>=4
L10 256861 SEA ABB=ON PLU=ON C5-C6-C6-C6/ES
L11 STR L2
L12 35 SEA SUB=L10 CSS SAM L11 AND L3 NOT L4
L13 0 SEA SUB=L10 CSS SAM L2 AND L3 NOT L4
L14 4008 SEA SUB=L10 CSS FUL L11 AND L3 NOT L4
D QUE L2
L15 4 SEA SUB=L14 SSS SAM L2
D SCA
L16 106 SEA SUB=L14 SSS FUL L2
SAV TEM L14 AUD807F0/A
SAV TEM L16 AUD807S0/A

FILE 'HCAPLUS' ENTERED AT 08:44:01 ON 24 JUN 2005

E MORRISON J/AU
L17 246 SEA ABB=ON PLU=ON ("MORRISON J"/AU OR "MORRISON J D"/AU)
E MORRISON JAMES/AU
L18 22 SEA ABB=ON PLU=ON ("MORRISON JAMES"/AU OR "MORRISON JAMES
DUNCAN"/AU)
E MORRISON JIM/AU
L19 4 SEA ABB=ON PLU=ON "MORRISON JIM"/AU
E LUCAS M/AU
L20 190 SEA ABB=ON PLU=ON ("LUCAS M"/AU OR "LUCAS M L"/AU)
E LUCAS MIKE/AU
E LUCAS MICHAEL/AU
L21 17 SEA ABB=ON PLU=ON ("LUCAS MICHAEL"/AU OR "LUCAS MICHAEL
L"/AU OR "LUCAS MICHAEL LESLIE"/AU)
E WHEELER S/AU
L22 56 SEA ABB=ON PLU=ON ("WHEELER S"/AU OR "WHEELER S A"/AU OR
"WHEELER S C"/AU OR "WHEELER S E"/AU OR "WHEELER S F"/AU OR
"WHEELER S G"/AU OR "WHEELER S H"/AU OR "WHEELER S J"/AU OR
"WHEELER S JAMES"/AU OR "WHEELER S L"/AU OR "WHEELER S M"/AU
OR "WHEELER S R"/AU OR "WHEELER S S"/AU OR "WHEELER S T"/AU OR
"WHEELER S V"/AU)
E WHEELER SARAH/AU
L23 16 SEA ABB=ON PLU=ON ("WHEELER SARAH"/AU OR "WHEELER SARAH
C"/AU OR "WHEELER SARAH CAROLINE"/AU OR "WHEELER SARAH E"/AU
OR "WHEELER SARAH J"/AU OR "WHEELER SARAH L"/AU)
L24 90 SEA ABB=ON PLU=ON L16
L25 QUE ABB=ON PLU=ON DRUG DELIVERY SYSTEMS+OLD,NT/CT
L26 0 SEA ABB=ON PLU=ON L24 AND L25

Search done by Noble Jarrell

L27 8 SEA ABB=ON PLU=ON L24 AND ?CONJUGAT?
 L28 0 SEA ABB=ON PLU=ON L16/D
 L29 0 SEA ABB=ON PLU=ON L24 AND (L17 OR L18 OR L19 OR L20 OR L21 OR L22)
 L30 5 SEA ABB=ON PLU=ON L14 AND (L17 OR L18 OR L19 OR L20 OR L21 OR L22)
 L31 17638 SEA ABB=ON PLU=ON L14 NOT L30
 L32 2551 SEA ABB=ON PLU=ON L31 AND ?CONJUGAT?
 L33 134 SEA ABB=ON PLU=ON L32 AND L25
 L34 QUE ABB=ON PLU=ON PY<=1999 OR AY<=1999 OR PRY<=1999 OR PD<19990730 OR AD<19990730 OR PRD<19990730
 L35 64 SEA ABB=ON PLU=ON L33 AND L34
 L36 7 SEA ABB=ON PLU=ON L27 AND L34
 L37 8 SEA ABB=ON PLU=ON L27 OR L36
 SEL AN L35 4 5 11 16 18 20-21 25 29 31-33 43-44 48 50
 L38 16 SEA ABB=ON PLU=ON ("119:146497"/AN OR "120:173477"/AN OR "121:286635"/AN OR "123:322102"/AN OR "126:308684"/AN OR "126:347323"/AN OR "127:99659"/AN OR "128:286354"/AN OR "130:213559"/AN OR "131:314185"/AN OR "132:26813"/AN OR "132:313463"/AN OR "132:73648"/AN OR "133:140227"/AN OR "136:107571"/AN OR "136:665"/AN OR "1993:546497"/AN OR "1994:173477"/AN OR "1994:686635"/AN OR "1995:721131"/AN OR "1997:218959"/AN OR "1997:372273"/AN OR "1997:463447"/AN OR "1998:208387"/AN OR "1999:185918"/AN OR "1999:708452"/AN OR "1999:722480"/AN OR "1999:764076"/AN OR "2000:10612"/AN OR "2000:508917"/AN OR "2001:844884"/AN OR "2002:72799"/AN) AND L35
 SEL AN 1 2 5 8 16 L38
 L39 11 SEA ABB=ON PLU=ON L38 NOT ("119:146497"/AN OR "130:213559"/AN OR "132:26813"/AN OR "136:107571"/AN OR "136:665"/AN OR "1993:546497"/AN OR "1999:185918"/AN OR "1999:764076"/AN OR "2001:844884"/AN OR "2002:72799"/AN)
 L40 19 SEA ABB=ON PLU=ON L37 OR L39

 FILE 'EMBASE' ENTERED AT 09:22:40 ON 24 JUN 2005
 L41 0 SEA ABB=ON PLU=ON L16
 L42 11483 SEA ABB=ON PLU=ON L14
 L43 1146 SEA ABB=ON PLU=ON L42 AND ?CONJUGAT?
 E ORAL/CT
 E E5+ALL
 E E2+ALL
 L44 109 SEA ABB=ON PLU=ON ORAL DRUG ADMINISTRATION/CT AND L43
 E MORRISON J/AU
 L45 338 SEA ABB=ON PLU=ON ("MORRISON J"/AU OR "MORRISON J D"/AU)
 E MORRISON JIM/AU
 E MORRISON JAMES/AU
 E LUCAS M/AU
 L46 326 SEA ABB=ON PLU=ON ("LUCAS M"/AU OR "LUCAS M L"/AU)
 E LUCAS MICHAEL/AU
 E WHEELER S/AU
 L47 151 SEA ABB=ON PLU=ON ("WHEELER S"/AU OR "WHEELER S A"/AU OR "WHEELER S B"/AU OR "WHEELER S C"/AU OR "WHEELER S D"/AU OR "WHEELER S E"/AU OR "WHEELER S F"/AU OR "WHEELER S G"/AU OR "WHEELER S H"/AU OR "WHEELER S J"/AU OR "WHEELER S K"/AU OR "WHEELER S L"/AU OR "WHEELER S M"/AU OR "WHEELER S R"/AU OR "WHEELER S V"/AU OR "WHEELER S W"/AU)
 L48 4 SEA ABB=ON PLU=ON L42 AND (L45 OR L46 OR L47)
 L49 109 SEA ABB=ON PLU=ON L44 NOT L48
 L50 91 SEA ABB=ON PLU=ON L49 AND PY<=1999

FILE 'EMBASE' ENTERED AT 09:33:50 ON 24 JUN 2005

SEL AN L50 60 63 40 43 46 12 16 20 29 2 3 5
 L51 12 SEA ABB=ON PLU=ON (1999046272/AN OR 1999256785/AN OR 1999324376/AN OR 84147710/AN OR 85080967/AN OR 88276787/AN OR 89150062/AN OR 90226900/AN OR 92292458/AN OR 94299511/AN OR 95240331/AN OR 96132330/AN) AND L50

```

FILE 'BIOSIS' ENTERED AT 09:34:08 ON 24 JUN 2005
L52      9899 SEA ABB=ON  PLU=ON  L14 OR L16
L53      1104 SEA ABB=ON  PLU=ON  L52 AND ?CONJUGAT?
          E MORRISON J/AU
L54      392 SEA ABB=ON  PLU=ON  ("MORRISON J"/AU OR "MORRISON J D"/AU)
          E MORRISON JAMES/AU
L55      4 SEA ABB=ON  PLU=ON  ("MORRISON JAMES"/AU OR "MORRISON JAMES
          D"/AU)
          E M LUCAS M/AU
          E LUCAS M/AU
L56      280 SEA ABB=ON  PLU=ON  ("LUCAS M"/AU OR "LUCAS M L"/AU)
          E LUCAS MICHAEL/AU
L57      3 SEA ABB=ON  PLU=ON  ("LUCAS MICHAEL"/AU OR "LUCAS MICHAEL
          L"/AU)
          E WHEELER S/AU
L58      160 SEA ABB=ON  PLU=ON  ("WHEELER S"/AU OR "WHEELER S A"/AU OR
          "WHEELER S C"/AU OR "WHEELER S CHRISTIAN"/AU OR "WHEELER S
          D"/AU OR "WHEELER S E"/AU OR "WHEELER S F"/AU OR "WHEELER S
          G"/AU OR "WHEELER S G B"/AU OR "WHEELER S H"/AU OR "WHEELER S
          J"/AU OR "WHEELER S JAMES"/AU OR "WHEELER S K"/AU OR "WHEELER
          S L"/AU OR "WHEELER S M"/AU OR "WHEELER S P"/AU OR "WHEELER S
          R"/AU OR "WHEELER S V"/AU OR "WHEELER S W"/AU)
          E WHEELER SARA/AU
L59      21 SEA ABB=ON  PLU=ON  ("WHEELER SARAH"/AU OR "WHEELER SARAH
          C"/AU OR "WHEELER SARAH E"/AU OR "WHEELER SARAH L"/AU OR
          "WHEELER SCHILLING T"/AU)
L60      1 SEA ABB=ON  PLU=ON  L53 AND (L54 OR L55 OR L56 OR L57 OR L58
          OR L59)

```

=> b reg

FILE 'REGISTRY' ENTERED AT 09:38:18 ON 24 JUN 2005
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
 provided by InfoChem.

STRUCTURE FILE UPDATES: 23 JUN 2005 HIGHEST RN 852898-09-0
 DICTIONARY FILE UPDATES: 23 JUN 2005 HIGHEST RN 852898-09-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when
 conducting SmartSELECT searches.

```

*****
*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*
*****

```


Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more
 information enter HELP PROP at an arrow prompt in the file or refer
 to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=> d que sta l14

Search done by Noble Jarrell

L3 SCR 1841
 L4 SCR 2039 OR 2041 OR 2050 OR 2049 OR 2053 OR 2052 OR 2048 O
 R 2043 OR 2054
 L10 256861 SEA FILE=REGISTRY ABB=ON PLU=ON C5-C6-C6-C6/ES
 L11 STR



VAR G1=C/21/23
 NODE ATTRIBUTES:
 CONNECT IS M1 RC AT 19
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED


GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 24

STEREO ATTRIBUTES: NONE
 L14 4008 SEA FILE=REGISTRY SUB=L10 CSS FUL L11 AND L3 NOT L4

100.0% PROCESSED 201370 ITERATIONS
 SEARCH TIME: 00.00.10

4008 ANSWERS

=> d que sta l16
 L2 STR




VAR G1=C/21/23
 NODE ATTRIBUTES:
 CONNECT IS M1 RC AT 19
 DEFAULT MLEVEL IS ATOM
 GGCAT IS UNS AT 18
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 24

STEREO ATTRIBUTES: NONE

L3 SCR 1841
L4 SCR 2039 OR 2041 OR 2050 OR 2049 OR 2053 OR 2052 OR 2048 O
R 2043 OR 2054
L10 256861 SEA FILE=REGISTRY ABB=ON PLU=ON C5-C6-C6-C6/ES
L11 STR



VAR G1=C/21/23

NODE ATTRIBUTES:

CONNECT IS M1 RC AT 19
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 24

STEREO ATTRIBUTES: NONE

L14 4008 SEA FILE=REGISTRY SUB=L10 CSS FUL L11 AND L3 NOT L4
L16 106 SEA FILE=REGISTRY SUB=L14 SSS FUL L2

100.0% PROCESSED 4008 ITERATIONS
SEARCH TIME: 00.00.01

106 ANSWERS

=> b hcap

FILE 'HCAPLUS' ENTERED AT 09:38:45 ON 24 JUN 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 24 Jun 2005 VOL 143 ISS 1
FILE LAST UPDATED: 23 Jun 2005 (20050623/ED)

Search done by Noble Jarrell

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d all fhitstr 130 tot

L30 ANSWER 1 OF 5 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2004:394158 HCAPLUS
 DN 141:47626
 ED Entered STN: 14 May 2004
 TI Absorption of the cholic acid-conjugated peptide hormone cholylsecretin from the rat ileum in vivo
 AU McHarg, S.; Morton, J. S.; McGinn, B. J.; Yasin, M.; Morrison, J. D.
 CS Institute of Biomedical and Life Sciences, University of Glasgow, Glasgow, UK
 SO Acta Physiologica Scandinavica (2004), 181(1), 23-34
 CODEN: APSCAX; ISSN: 0001-6772
 PB Blackwell Publishing Ltd.
 DT Journal
 LA English
 CC 2-6 (Mammalian Hormones)
 AB Previously, the authors demonstrated that gastrin peptides as long as 34 amino acids were absorbed from the ileum of rat after conjugation to the C24 position of cholic acid and that these peptides retained full biol. activity. As absorption was specific to the ileum, it was inferred that the conjugated hormone was taken up by the bile salt transporters. The authors have now extended these expts. to a member of a different family of hormones, viz. secretin, a 27-amino acid hormone that stimulates serous secretions from the exocrine pancreas. After conjugation to cholic acid, the degree of cholylsecretin absorption from the ileum of anesthetized rats was assessed from the increase in pancreatic secretions. A complication to the study was that intra-ileal infusion of native secretin caused a transient increase in the levels of pancreatic secretions. This was in contrast to the effects of intra-ileal infusion of cholylsecretin which did not cause this transient increase but, instead, gave rise to a delayed increase in pancreatic secretions which was sustained over several hours during which cholylsecretin was detected in plasma in high concentration by mass spectrometry. The pancreatic response to cholylsecretin was abolished by co-infusion of 50 mM taurocholate, employed to compete with the bile salt transporters, although a transient increase in pancreatic secretions similar to that caused by secretin was now generated. This was shown to arise from an action of taurocholate per se causing the release of endogenous secretin which is present in rat ileum. The authors, therefore, concluded that cholylsecretin had been absorbed from the rat ileum by uptake by bile salt transporters.
 ST secretin cholic acid conjugate bile salt absorption ileum rat;
 cholylsecretin absorption bile salt transporter ileum
 IT Pancreas
 (absorption of cholic acid-conjugated peptide hormone cholylsecretin from the rat ileum in vivo)
 IT Transport proteins
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (bile salt; absorption of cholic acid-conjugated peptide hormone cholylsecretin from the rat ileum in vivo)
 IT Intestine
 (ileum; absorption of cholic acid-conjugated peptide hormone cholylsecretin from the rat ileum in vivo)
 IT Bile salts
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (transporter; absorption of cholic acid-conjugated peptide hormone cholylsecretin from the rat ileum in vivo)
 IT Biological transport
 (uptake; absorption of cholic acid-conjugated peptide hormone cholylsecretin from the rat ileum in vivo)

Search done by Noble Jarrell

IT 71-52-3, Bicarbonate, biological studies 709002-71-1
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (absorption of cholic acid-conjugated peptide hormone cholylsecretin
 from the rat ileum in vivo)

RE.CNT 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Baker, R; Proc Soc Exp Biol Med 1960, V105, P521 HCAPLUS
- (2) Bayliss, W; J Physiol 1902, V28, P325 HCAPLUS
- (3) Bodanszky, A; J Am Chem Soc 1969, V91, P944 HCAPLUS
- (4) Chey, W; J Physiol 1982, V324, P263 HCAPLUS
- (5) Dakka, T; Digestion 1995, V56, P165 HCAPLUS
- (6) Dietschy, J; J Lipid Res 1968, V9, P297 HCAPLUS
- (7) Gossen, D; Biochem Biophys Res Commun 1989, V160, P862 HCAPLUS
- (8) Gourlet, P; Eur J Biochem 1996, V239, P349 HCAPLUS
- (9) Gourlet, P; Peptides 1996, V17, P825 HCAPLUS
- (10) Gronenborn, A; FEBS Lett 1987, V215, P88 HCAPLUS
- (11) Horvath, K; J Assoc Acad Min Phys 1998, V9, P9 MEDLINE
- (12) Izzo, R; Int J Pept Prot Res 1984, V23, P292 HCAPLUS
- (13) Jin, H; Gastroenterology 1993, V105, P208 HCAPLUS
- (14) Kern, J; J Autism Develop Disord 2002, V32, P153
- (15) Larsson, L; Histochemistry 1978, V58, P23 HCAPLUS
- (16) Morely, J; Nature 1965, V207, P1356
- (17) Moriyasu, M; Pancreas 1994, V9, P129 MEDLINE
- (18) Morrison, J; J Physiol Proceedings 2004
- (19) New, R; Int Patent Appl WO 96/06635 1996
- (20) O'Maille, E; J Physiol 1965, V180, P67 HCAPLUS
- (21) Playoust, M; J Clin Invest 1964, V43, P467 HCAPLUS
- (22) Roberts, P; Digestion 1999, V60, P332 HCAPLUS
- (23) Straus, E; Gastroenterology 1978, V75, P401 HCAPLUS
- (24) The Diabetes Control And Complications Trial Research Group; New Engl J Med 2000, V342, P381
- (25) Tranberg, K; Ann Surg 1985, V201, P300 HCAPLUS
- (26) Weiner, I; Handbook of Physiology - Alimentary Canal III 1967, P1439
- (27) Wettergren, A; Digest Dis Sci 1993, V38, P665 HCAPLUS
- (28) Wheeler, S; Acta Physiol Scand 2002, V176, P203 HCAPLUS
- (29) Wheeler, S; Exp Physiol 1997, V82, P729 HCAPLUS

IT 709002-71-1


RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (absorption of cholic acid-conjugated peptide hormone cholylsecretin
 from the rat ileum in vivo)

RN 709002-71-1 HCAPLUS


CN L-Valinamide, N-[(3 α , 5 β , 7 α , 12 α)-3,7,12-trihydroxy-
 24-oxocholan-24-yl]-L-histidyl-L-seryl-L- α -aspartylglycyl-L-threonyl-
 L-phenylalanyl-L-threonyl-L-seryl-L- α -glutamyl-L-leucyl-L-seryl-L-
 arginyl-L-leucyl-L-arginyl-L- α -glutamylglycyl-L-alanyl-L-arginyl-L-
 leucyl-L-glutamyl-L-arginyl-L-leucyl-L-leucyl-L-glutamylglycyl-L-
 leucyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A





PAGE 1-B




PAGE 1-C



PAGE 2-A





PAGE 3-A




PAGE 3-B

Bu-i

PAGE 4-A



PAGE 4-B



L30 ANSWER 2 OF 5 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2002:869795 HCAPLUS
 DN 138:181158
 ED Entered STN: 17 Nov 2002
 TI Absorption of biologically active peptide hormones from the small intestine of rat
 AU Wheeler, S.; McGinn, B. J.; Lucas, M. L.; Morrison, J. D.
 CS University of Glasgow, Glasgow, G12 8QQ, UK
 SO Acta Physiologica Scandinavica (2002), 176(3), 203-213
 CODEN: APSCAX; ISSN: 0001-6772
 PB Blackwell Science Ltd.
 DT Journal
 LA English
 CC 2-6 (Mammalian Hormones)
 AB Absorption of the 4, 10 and 34 amino acid forms of gastrin from the small intestine has been investigated in anesthetized rats. The method of assessment of successful absorption of the hormone into the systemic circulation was when the amount of acid secreted by the stomach over consecutive 15-min periods was increased. When the natural hormones were infused into the ileum in a relatively high dose, there was no increase in gastric acid secretion, indicating that they had not been absorbed. Each of the forms of gastrin was conjugated at the free N-terminus to the carboxyl group of cholic acid. Subsequent infusion of the conjugated form of gastrin into the ileum, this time in relatively low doses, resulted in substantial and prolonged increases in gastric acid secretion, indicating that these hormones had been successfully absorbed. In addition, conjugation of the 10 and 34 amino acid forms of gastrin with cholic acid was shown to increase markedly the potency in evoking an increase in gastric acid secretion in response to i.v. injection of the hormone. Absorption of the gastrin conjugates was specific to the ileum thus indicating that they had been absorbed through the bile salt transporters.
 ST gastrin isoform absorption small intestine rat
 IT Transport proteins
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (bile salt; absorption of biol. active peptide hormones from ileum of rat indicates absorption through bile salt transporters)
 IT Intestine (ileum; absorption of biol. active peptide hormones from the small intestine of rat)
 IT Gastric acid (secretion; absorption of biol. active peptide hormones from the small intestine of rat)
 IT Circulation (systemic; absorption of biol. active peptide hormones from the small intestine of rat)
 IT Bile salts
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (transporter; absorption of biol. active peptide hormones from ileum of rat indicates absorption through bile salt transporters)
 IT Biological transport (uptake; absorption of biol. active peptide hormones from the small intestine of rat)
 IT 1947-37-1, 4-7-Cholecystokinin-7 (swine) 18828-47-2 171511-54-9 324753-46-0 496946-81-7 499210-69-4 499210-82-1
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (absorption of biol. active peptide hormones from the small intestine of rat)
 IT 81-25-4, Cholic acid
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (conjugate; absorption of biol. active peptide hormones from the small intestine of rat)
 RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Baker, R; Proc Soc Exp Biol Med 1960, V105, P521 HCAPLUS

- (2) Bendayan, M; Diabetologica 1990, V33, P197 HCAPLUS
- (3) Burkoth, T; Crit Rev Therap Drug Carr Sys 1999, V16, P331 HCAPLUS
- (4) Chan, W; Fmoc Solid Phase Peptide Synthesis -- a Practical Approach 2000
- (5) Dakka, T; Digestion 1995, V56, P165 HCAPLUS
- (6) Dange, C; J Pharmaceut Sci 1997, V86, P1403 HCAPLUS
- (7) Davenport, H; Physiology of the Digestive Tract, 4th edn 1977
- (8) Haas, S; Vaccine 1996, V14, P785 HCAPLUS
- (9) Holt, P; Am J Physiol 1964, V207, P1 HCAPLUS
- (10) Kramer, W; US 5462933 1995 HCAPLUS
- (11) Kramer, W; J Biol Chem 1994, V269, P10621 HCAPLUS
- (12) Krchnak, V; Int J Pept Prot Res 1988, V32, P415 HCAPLUS
- (13) Lack, L; Am J Physiol 1961, V200, P313 HCAPLUS
- (14) Lack, L; Am J Physiol 1966, V210, P1142 HCAPLUS
- (15) Layer, P; Digest Dis Sci 1995, V40, P1074 HCAPLUS
- (16) Lichtberger, L; Gastroenterology 1986, V90, P1223
- (17) Mathlowitz, E; Nature 1997, V386, P410
- (18) Matthews, D; Gastroenterology 1976, V71, P151 HCAPLUS
- (19) McGinn, B; Proteins Labfax 1996, P139 HCAPLUS
- (20) Morley, J; Nature 1965, V207, P1356 HCAPLUS
- (21) Muranishi, S; J Contr Release 1992, V19, P179 HCAPLUS
- (22) Namba, M; Regul Pept 1986, V15, P121 HCAPLUS
- (23) Playoust, M; J Clin Invest 1964, V43, P467 HCAPLUS
- (24) Ritschel, W; Meth Find Exp Clin Pharmacol 1991, V13, P205 HCAPLUS
- (25) Roberts, P; Digestion 1999, V60, P332 HCAPLUS
- (26) Royal Pharmaceutical Society; British National Formulary 2001
- (27) Skyler, J; Lancet 2001, V357, P331 HCAPLUS
- (28) Stephan, Z; Biochem Pharmacol 1992, V43, P1969 HCAPLUS
- (29) Swaan, P; Adv Drug Del Rev 1996, V20, P59 HCAPLUS
- (30) Swaan, P; Bioconj Chem 1997, V8, P520 HCAPLUS
- (31) Trier, J; Gastroenterol Clin N Am 1991, V20, P531 MEDLINE
- (32) Walsh, J; Gut Peptides: Biochemistry and Physiology 1994, P75 HCAPLUS
- (33) Weiner, I; Handbook of Physiology -- Alimentary Canal 1967, VIII, P1439
- (34) Wheeler, S; Exp Physiol 1997, V82, P729 HCAPLUS

IT 171511-54-9


RL: BSU (Biological study, unclassified); BIOL (Biological study)
(absorption of biol. active peptide hormones from the small intestine
of rat)

RN 171511-54-9 HCAPLUS


CN L-Phenylalaninamide, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-
trihydroxy-24-oxocholan-24-yl]-L-tryptophyl-L-methionyl-L- α -aspartyl-
(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

PAGE 1-A



PAGE 2-A



L30 ANSWER 3 OF 5 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2001:101167 HCAPLUS
 DN 134:168315
 ED Entered STN: 09 Feb 2001
 TI Enhancement of bioavailability of peptides with bile salts
 IN Morrison, James Duncan; Lucas, Michael Leslie;
 Wheeler, Sarah
 PA The University Court of the University of Glasgow, UK
 SO PCT Int. Appl., 28 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM C07J
 CC 63-5 (Pharmaceuticals)
 Section cross-reference(s): 1
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001009163	A2	20010208	WO 2000-GB2903	20000728
WO 2001009163	A3	20010907		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,

Search done by Noble Jarrell

CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
 HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,
 LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU,
 SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,
 YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
 CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

GB 2355009 A1 20010411 GB 1999-17793 19990730
 AU 2000061739 A5 20010219 AU 2000-61739 20000728
 EP 1228093 A2 20020807 EP 2000-948177 20000728

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL

PRAI GB 1999-17793 A 19990730
 WO 2000-GB2903 W 20000728

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 2001009163	ICM	C07J
WO 2001009163	ECLA	A61K047/48H4; C07K014/47; C07K014/575; C07K014/595
GB 2355009	ECLA	A61K047/48H4; C07K014/47; C07K014/575; C07K014/595
OS	MARPAT 134:168315	
AB	The present invention relates to improving and/or increasing the bioavailability of a biol. active substance, such as a peptide. In particular the present invention relates to the conjugation of the biol. active substance to a bile acid. The conjugated biol. active substance is suitable particularly for oral or parental administration. Ileal administration of 600µg/kg gastrin tetrapeptide conjugated to cholate resulted in a significant mean increase in gastric acid secretion of 1.84 µmol over a 3 h collection period, while no increase in acid secretion was noticed by administration of tetragastrin alone or with sep. cholate.	
ST	bioavailability enhancement peptide bile salt	
IT	Immunoglobulins RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (A; enhancement of bioavailability of peptides with bile salts)	
IT	Immunoglobulins RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (D; enhancement of bioavailability of peptides with bile salts)	
IT	Immunoglobulins RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (E; enhancement of bioavailability of peptides with bile salts)	
IT	Immunoglobulins RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (G; enhancement of bioavailability of peptides with bile salts)	
IT	Immunoglobulins RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (M; enhancement of bioavailability of peptides with bile salts)	
IT	Chemotherapy (agents; enhancement of bioavailability of peptides with bile salts)	
IT	Adrenoceptors RL: BSU (Biological study, unclassified); BIOL (Biological study) (antagonists; enhancement of bioavailability of peptides with bile salts)	
IT	Anemia (disease) (antianemic factors; enhancement of bioavailability of peptides with bile salts)	

- IT Peptides, biological studies
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)
 (conjugates; enhancement of bioavailability of peptides with bile salts)
- IT Adrenoceptor agonists
 Adrenoceptor antagonists
 Analgesics
 Anesthetics
 Anti-inflammatory agents
 Antianginal agents
 Antiarrhythmics
 Antibacterial agents
 Anticoagulants
 Anticonvulsants
 Antidepressants
 Antihistamines
 Antiparkinsonian agents
 Antipsychotics
 Antiviral agents
 Anxiolytics
 Cardiotonics
 Diuretics
 Drug bioavailability
 Fungicides
 Hypnotics and Sedatives
 Hypolipemic agents
 Muscarinic agonists
 Muscarinic antagonists
 Nicotinic antagonists
 Parasitocides
 Permeation enhancers
 Stomach
 Vasodilators
 (enhancement of bioavailability of peptides with bile salts)
- IT Peptides, biological studies
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
 (enhancement of bioavailability of peptides with bile salts)
- IT Antibodies
 Blood-coagulation factors
 Ferritins
 Glycoproteins, general, biological studies
 Hemoglobins
 Interferons
 Oligonucleotides
 Opioids
 Polynucleotides
 Polysaccharides, biological studies
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (enhancement of bioavailability of peptides with bile salts)
- IT Bile acids
 RL: BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
 (enhancement of bioavailability of peptides with bile salts)
- IT Bile salts
 RL: BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
 (enhancement of bioavailability of peptides with bile salts)
- IT Gastrointestinal motility
 (gastric, drugs for treatment of; enhancement of bioavailability of

- peptides with bile salts)
- IT Drug delivery systems
(oral; enhancement of bioavailability of peptides with bile salts)
- IT Drug delivery systems
(parenterals; enhancement of bioavailability of peptides with bile salts)
- IT Antiulcer agents
(peptic; enhancement of bioavailability of peptides with bile salts)
- IT Neuropeptides
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(transmitters; enhancement of bioavailability of peptides with bile salts)
- IT 9001-08-5D, inhibitor
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(anticholinesterase; enhancement of bioavailability of peptides with bile salts)
- IT 50-56-6, Oxytocin, biological studies 1393-25-5, Secretin 8001-27-2, Hirudin 9001-05-2, Catalase 9001-27-8, Factor viii 9001-28-9, Factor IX 9002-60-2, Acth, biological studies 9002-61-3, Chorionic gonadotropin 9002-64-6, Parathyroid hormone 9002-67-9, Luteinizing hormone 9002-68-0, Follicle stimulating hormone 9002-71-5, Thyroid stimulating hormone 9002-72-6, Somatotropin 9002-76-0, Gastrin 9004-10-8, Insulin, biological studies 9007-12-9, Calcitonin 9007-43-6, Cytochrome c, biological studies 9007-92-5, Glucagon, biological studies 9011-97-6, Cholecystokinin 9015-71-8, Corticotropin releasing hormone 9015-94-5, Renin, biological studies 9034-39-3, Growth hormone releasing hormone 9034-40-6, Gonadotropin releasing hormone 9038-70-4, Somatomedin 9039-53-6, Urokinase 9041-90-1, Angiotensin I 9054-89-1, Superoxide dismutase 9087-70-1, Aprotinin 11000-17-2, Antidiuretic hormone 11096-26-7, Erythropoietin 11128-99-7, Angiotensin II 24305-27-9, Thyrotropin releasing hormone 51110-01-1, Somatostatin 57285-09-3, Inhibin 59392-49-3, Gastric inhibitory peptide 67763-96-6, Igf1 67763-97-7, Igf2 80043-53-4, Gastrinreleasing peptide 85637-73-6, Atrial natriuretic hormone 89750-14-1, Glucagon-like peptide I 119418-04-1, Galanin 139639-23-9, Tissue plasminogen activator
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(enhancement of bioavailability of peptides with bile salts)
- IT 79-14-1D, Glycolic acid, salts 81-24-3D, Taurocholic acid, salts 81-25-4, Cholic acid 83-44-3D, Deoxycholic acid, salts 128-13-2D, Ursodeoxycholic acid, salts 360-65-6D, Glycodeoxycholic acid, salts 474-25-9D, Chenodeoxycholic acid, salts 474-74-8D, Glycolithocholic acid, salts 516-35-8D, Taurochenodeoxycholic acid, salts 516-50-7D, Taurodeoxycholic acid, salts 516-90-5D, TAurolithocholic acid, salts 640-79-9D, Glycochenodeoxycholic acid, salts 14605-22-2D, Tauroursodeoxycholic acid, salts 63948-32-3 64480-66-6D, Glycoursodeoxycholic acid, salts 83381-47-9, Gastrin-34 I (rat) 171511-54-9 324753-46-0 325142-35-6
RL: BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(enhancement of bioavailability of peptides with bile salts)
- IT 9003-99-0, Peroxidase
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(horseradish; enhancement of bioavailability of peptides with bile salts)
- IT 9002-10-2, Tyrosinase
RL: BAC (Biological activity or effector, except adverse); BSU (Biological

study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(mushroom; enhancement of bioavailability of peptides with bile salts)

IT 9035-81-8, Trypsin inhibitor

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(soy bean; enhancement of bioavailability of peptides with bile salts)


IT 81-25-4, Cholic acid

RL: BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(enhancement of bioavailability of peptides with bile salts)

RN 81-25-4 HCAPLUS

CN Cholan-24-oic acid, 3,7,12-trihydroxy-, (3 α ,5 β ,7 α ,12 α a.)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L30 ANSWER 4 OF 5 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1991:580392 HCAPLUS

DN 115:180392

ED Entered STN: 01 Nov 1991

TI The effect of sodium deoxycholate and other surfactants on the mucosal surface pH in proximal jejunum of rat

AU McKie, A. T.; Stewart, W.; Lucas, M. L.

CS Inst. Physiol., Glasgow Univ., Glasgow, G12 8QQ, UK

SO Naunyn-Schmiedeberg's Archives of Pharmacology (1991), 343(6), 659-64

CODEN: NSAPCC; ISSN: 0028-1298

DT Journal

LA English

CC 13-7 (Mammalian Biochemistry)


AB The mucosal surface pH (acid microclimate) and nucleotide levels of rat proximal jejunum were measured in vivo under various conditions which included exposure to pharmacol. agents and to surfactants. Mucosal surface pH was unaffected by sodium nitroprusside, A 23187, and amiloride, as was mucosal cGMP content, although amiloride and A 23187 reduced cAMP content. In contrast, surfactants elevated the pH of the mucosal surface significantly: control value 6.23; Lubrol PX 0.8% (volume/volume) 6.98 sodium deoxycholate 2 mM 6.67; Triton X 100 0.5% (volume/volume) 7.41. No significant changes in cGMP levels were noted after surfactant treatment, although deoxycholate and Triton X 100 reduced cAMP levels. The ability of higher concns. of surfactant to elevate the mucosal surface pH beyond neutrality to values similar to plasma pH contrasts with the action of Escherichia coli heat-stable (STa) enterotoxin, which at high concns. could not elevate the mucosal surface pH beyond neutrality. Consistent with the known effects on tight junction permeability, surfactants may act by allowing plasma-like subepithelial fluid to neutralize the microclimate.

ST jejunum mucosa pH surfactant; cAMP jejunum mucosa pH surfactant; cGMP jejunum mucosa pH surfactant

IT Surfactants

- (proximal jejunum mucosal surface pH response to, ion movements and cyclic nucleotides in relation to)
- IT Intestine
(jejunum, proximal, mucosa, surface pH of, surfactants effect on, ion movements and cyclic nucleotides in relation to)
- IT 7440-23-5, Sodium, biological studies
RL: BIOL (Biological study)
(hydrogen ion exchange with, in jejunum mucosa, effect of surfactant on surface pH in relation to)
- IT 60-92-4, CAMP 7665-99-8, CGMP
RL: BIOL (Biological study)
(of jejunum mucosa, surfactants effect on, mucosal surface pH in relation to)
- IT 302-95-4, Sodium deoxycholate 577-11-7 9002-92-0, Lubrol PX
9002-93-1, Triton X-100
RL: BIOL (Biological study)
(proximal jejunum mucosal surface pH response to, ion movements and cyclic nucleotides in relation to)
- IT 12408-02-5, Hydrogen ion, biological studies
RL: BIOL (Biological study)
(sodium exchange with, in jejunum mucosa, effect of surfactant on surface pH in relation to)
- IT 7440-70-2, Calcium, biological studies
RL: BIOL (Biological study)
(transport of, by jejunum mucosa, surfactants effect on surface pH and cyclic nucleotides in relation to)
- IT 302-95-4, Sodium deoxycholate
RL: BIOL (Biological study)
(proximal jejunum mucosal surface pH response to, ion movements and cyclic nucleotides in relation to)
- RN 302-95-4 HCAPLUS
- CN Cholan-24-oic acid, 3,12-dihydroxy-, monosodium salt,
(3 α ,5 β ,12 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● Na


L30 ANSWER 5 OF 5 HCAPLUS COPYRIGHT 2005 ACS on STN
AN 1984:32553 HCAPLUS
DN 100:32553
ED Entered STN: 12 May 1984
TI The effect of deoxycholate on intestinal surface pH and
5-methyltetrahydropteroylglutamate absorption in the rat proximal jejunum
in vitro
AU Blair, John A.; Hilburn, Michael E.; Lucas, Michael L.; Said,
Hamid M.
CS Dep. Chem., Univ. Aston, Birmingham, B4 7ET, UK
SO Biochemical Society Transactions (1983), 11(2), 165-7

Search done by Noble Jarrell

CODEN: BCSTB5; ISSN: 0300-5127

DT Journal
 LA English
 CC 13-2 (Mammalian Biochemistry)
 AB The effects of deoxycholate (0.01-10 mM) on rat proximal jejunum in vitro indicated that intestinal surface pH is a determinant of folate absorption.
 ST intestine pH folate absorption deoxycholate
 IT Bile acids
 RL: BIOL (Biological study)
 (folate absorption by intestinal jejunum response to, surface pH in relation to)
 IT Intestine, metabolism
 (proximal jejunum, folate absorption by, deoxycholate effect on, surface pH in relation to)
 IT 59-30-3, biological studies 134-35-0
 RL: BIOL (Biological study)
 (absorption of, by proximal jejunum, deoxycholate effect on, surface pH in relation to)
 IT 81-25-4 360-65-6
 RL: BIOL (Biological study)
 (folate absorption by intestinal jejunum response to, surface pH in relation to)
 IT 83-44-3
 RL: BIOL (Biological study)
 (folate absorption by proximal jejunum response to, surface pH in relation to)
 IT 81-25-4
 RL: BIOL (Biological study)
 (folate absorption by intestinal jejunum response to, surface pH in relation to)
 RN 81-25-4 HCAPLUS
 CN Cholan-24-oic acid, 3,7,12-trihydroxy-, (3 α ,5 β ,7 α ,12 α -a.)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> d all hitstr 140 tot

L40 ANSWER 1 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2002:537280 HCAPLUS
 DN 138:85395
 ED Entered STN: 19 Jul 2002
 TI Participation of two members of the very long-chain acyl-CoA synthetase family in bile acid synthesis and recycling
 AU Mihalik, Stephanie J.; Steinberg, Steven J.; Pei, Zhengtong; Park, Joseph; Kim, Do G.; Heinzer, Ann K.; Dacremont, Georges; Wanders, Ronald J. A.; Cuebas, Dean A.; Smith, Kirby D.; Watkins, Paul A.
 CS Kennedy Krieger Institute and the Department of Pediatrics, Johns Hopkins

Search done by Noble Jarrell

University School of Medicine, Baltimore, MD, 21205, USA

SO Journal of Biological Chemistry (2002), 277(27), 24771-24779
CODEN: JBCHA3; ISSN: 0021-9258

PB American Society for Biochemistry and Molecular Biology

DT Journal

LA English

CC 7-2 (Enzymes)
Section cross-reference(s): 3, 13

OS CASREACT 138:85395

AB Bile acids are synthesized de novo in the liver from cholesterol and conjugated to glycine or taurine via a complex series of reactions involving multiple organelles. Bile acids secreted into the small intestine are efficiently reabsorbed and reutilized. Activation by thioesterification to CoA is required at two points in bile acid metabolism. First, 3 α ,7 α ,12 α -trihydroxy-5 β -cholestanoic acid, the 27-carbon precursor of cholic acid, must be activated to its CoA derivative before side chain cleavage via peroxisomal β -oxidation. Second, reutilization of cholate and other C24 bile acids requires reactivation prior to re-conjugation. We reported previously that homolog 2 of very long-chain acyl-CoA synthetase (VLCS) can activate cholate. We now show that homolog 2 also activates chenodeoxycholate, the secondary bile acids deoxycholate and lithocholate, and 3 α ,7 α ,12 α -trihydroxy-5 β -cholestanoic acid. In contrast, VLCS activated 3 α ,7 α ,12 α -trihydroxy-5 β -cholestanoate, but did not utilize any of the C24 bile acids as substrates. We hypothesize that the primary function of homolog 2 is in the reactivation and recycling of C24 bile acids, whereas VLCS participates in the de novo synthesis pathway. Results of in situ hybridization, topog. orientation, and inhibition studies are consistent with the proposed roles of these enzymes in bile acid metabolism

ST very long chain acyl CoA synthetase bile acid recycling; mouse cDNA sequence bile acid CoA synthetase liver

IT Mus musculus
(VLCS cDNA sequence; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)

IT Bile acids
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(enzyme substrate specificity; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)

IT Liver
(hepatocyte, compartmentalized expression of VLCS; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)

IT Molecular topology
(membrane topol. of VLCS-H2; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)

IT Protein sequences
cDNA sequences
(of VLCS of mouse liver; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)

IT 114797-03-4P, Bile acid-CoA synthetase
RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PREP (Preparation)
(VLCS homolog 2; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)

- IT 69403-06-1P, Very long-chain acyl-CoA synthetase
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation)
 (VLCS; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- IT 470737-50-9P
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PREP (Preparation)
 (amino acid sequence; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- IT 83-44-3, Deoxycholic acid 434-13-9, Lithocholic acid 474-25-9, Chenodeoxycholic acid
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (enzyme substrate specificity; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- IT 547-98-8P
 RL: BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)
 (enzyme substrate specificity; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- IT 5226-26-6
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (in preparation of bile acid precursor THCA; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- IT 3396-82-5, Sodium cyanide (Na(14CN))
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (in preparation of 14C-labeled trihydroxycholestanate; bile acid-CoA synthetase activity and cellular localization of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- IT 114416-41-0P 114443-04-8P 114443-05-9P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (in preparation of 14C-labeled trihydroxycholestanate; bile acid-CoA synthetase activity and cellular localization of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- IT 200385-45-1, GenBank AF033031
 RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (nucleotide sequence; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- IT 114416-40-9P
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (14C-labeled enzyme substrate; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)
- RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Antonenkov, V; J Biol Chem 1997, V272, P26023 HCAPLUS
 (2) Bahar, R; Gastroenterol Clin North Am 1999, V28, P27 MEDLINE
 (3) Carey, M; The Liver: Biology and Pathobiology, 3rd Ed 1994, P719

- (4) Dieuaide-Noubhani, M; Eur J Biochem 1996, V240, P660 HCAPLUS
- (5) Giger, R; J Comp Neurol 1996, V375, P378 HCAPLUS
- (6) Hofmann, A; The Liver: Biology and Pathobiology, 3rd Ed 1994, P677
- (7) Kase, B; J Biol Chem 1989, V264, P9220 HCAPLUS
- (8) Kase, F; J Lipid Res 1983, V24, P1560 HCAPLUS
- (9) Katz, N; Eur J Biochem 1983, V135, P103 HCAPLUS
- (10) Katz, N; Eur J Biochem 1989, V180, P185 HCAPLUS
- (11) Killenberg, P; J Lipid Res 1978, V19, P24 HCAPLUS
- (12) Kurosawa, T; Anal Chim Acta 1998, V365, P249 HCAPLUS
- (13) Lim, W; Biochem J 1981, V197, P611 HCAPLUS
- (14) Lowry, O; J Biol Chem 1951, V193, P265 HCAPLUS
- (15) Polokoff, M; J Biol Chem 1977, V252, P1167 HCAPLUS
- (16) Rost, B; Protein Sci 1995, V4, P521 HCAPLUS
- (17) Schepers, L; Biochem J 1989, V257, P221 HCAPLUS
- (18) Setchell, K; Liver Disease in Children, 2nd Ed 2001, P701
- (19) Smith, B; Exp Cell Res 2000, V254, P309 HCAPLUS
- (20) Solaas, K; J Lipid Res 2000, V41, P1154 HCAPLUS
- (21) Steinberg, S; Ann Neurol 1999, V46, P409 HCAPLUS
- (22) Steinberg, S; Biochem Biophys Res Commun 1999, V257, P615 HCAPLUS
- (23) Steinberg, S; J Biol Chem 2000, V275, P15605 HCAPLUS
- (24) Steinberg, S; Mol Genet Metab 1999, V68, P32 HCAPLUS
- (25) Tserng, K; J Lipid Res 1977, V18, P400 HCAPLUS
- (26) Twisk, J; J Clin Invest 1995, V95, P1235 HCAPLUS
- (27) Uchiyama, A; J Biol Chem 1996, V271, P30360 HCAPLUS
- (28) van Grunsven, E; Proc Natl Acad Sci U S A 1998, V95, P2128 HCAPLUS
- (29) van Veldhoven, P; J Biol Chem 1992, V267, P20065 HCAPLUS
- (30) Vanhove, G; J Biol Chem 1993, V268, P10335 HCAPLUS
- (31) Vlahcevic, Z; Gastroenterol Clin North Am 1999, V28, P1 MEDLINE
- (32) Wanders, R; The Metabolic & Molecular Bases of Inherited Disease, 8th Ed 2001, P3219
- (33) Watkins, P; Am J Hum Genet 1995, V57, P292 HCAPLUS
- (34) Wheeler, J; Arch Biochem Biophys 1997, V348, P15 HCAPLUS
- (35) Xu, R; Biochem Biophys Res Commun 1996, V221, P271 HCAPLUS

IT 5226-26-6

RL: RCT (Reactant); RACT (Reactant or reagent)


(in preparation of bile acid precursor THCA; bile acid-CoA synthetase activity, cellular localization, and membrane topog. of two very long-chain acyl-CoA synthetase (VLCS) family proteins suggests a role in bile acid synthesis and recycling)

RN 5226-26-6 HCAPLUS

CN Cholest-24-en-26-oic acid, 3,7,12-trihydroxy-,
(3 α ,5 β ,7 α ,12 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry unknown.



L40 ANSWER 2 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2000:508917 HCAPLUS

DN 133:140227

ED Entered STN: 27 Jul 2000

TI Method and compositions for lipidization of hydrophilic molecules

IN Shen, Wei-chiang; Wang, Jinghua
 PA The University of Southern California, USA
 SO U.S., 34 pp.
 CODEN: USXXAM
 DT Patent
 LA English
 IC ICM A61K038-28
 INCL 514003000
 CC 63-6 (Pharmaceuticals)
 Section cross-reference(s): 1

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6093692	A	20000725	US 1997-936898	19970925 <--
PRAI	US 1996-77177P	P	19960926	<--	
	US 1997-49499P	P	19970613	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 6093692	ICM	A61K038-28
	INCL	514003000
US 6093692	NCL	514/003.000; 514/002.000; 514/009.000; 514/019.000; 514/023.000; 530/300.000; 530/303.000; 530/307.000; 530/315.000; 530/317.000; 530/331.000; 530/333.000; 530/350.000
	ECLA	A61K047/48H4 <--

OS MARPAT 133:140227

AB Fatty acid derivs. of disulfide-containing compds. (for example, disulfide-containing peptides or proteins) comprising fatty acid-conjugated products with a disulfide linkage are employed for delivery of the compds. to mammalian cells. This modification markedly increases the absorption of the compds. by mammalian cells relative to the rate of absorption of the unconjugated compds., as well as prolonging blood and tissue retention of the compds. Moreover, the disulfide linkage in the conjugate is quite labile in vivo and thus facilitates intracellular or extracellular release of the intact compds. from the fatty acid moieties. N-palmityl-2-pyridyldithiocysteine was prepared and reacted with Bowman-Birk inhibitor (BBI) to obtain a palmityl disulfide conjugate of BBI. When the conjugate was incubated with colon carcinoma cells (Caco-2) in serum-free medium, the uptake of the conjugate was higher than that of BBI.

ST fatty acid disulfide protein conjugate bioavailability

IT Antisense oligonucleotides

RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(complementary to mRNA of monoamine oxidase B, palmitylated;
 conjugates of hydrophilic mols. with fatty acid or steroid
 disulfide derivs. for improving their bioavailabilities)

IT Drug bioavailability

Drug delivery systems

(conjugates of hydrophilic mols. with fatty acid or steroid
 disulfide derivs. for improving their bioavailabilities)

IT Amino acids, biological studies

Carbohydrates, biological studies

Nucleosides, biological studies

Nucleotides, biological studies

Oligonucleotides

Peptides, biological studies

Proteins, general, biological studies

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(conjugates with fatty acid disulfide derivs.;

conjugates of hydrophilic mols. with fatty acid or steroid

disulfide derivs. for improving their bioavailabilities)

IT Fatty acids, biological studies

Steroids, biological studies

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)

- (disulfide derivs., conjugates with proteins;
conjugates of hydrophilic mols. with fatty acid or steroid
disulfide derivs. for improving their bioavailabilities)
- IT Biological transport
(drug; conjugates of hydrophilic mols. with fatty acid or
steroid disulfide derivs. for improving their bioavailabilities)
- IT Drug delivery systems
(liposomes; conjugates of hydrophilic mols. with fatty acid
or steroid disulfide derivs. for improving their bioavailabilities)
- IT 16679-58-6DP, Desmopressin, fatty acid disulfide conjugates
37330-34-ODP, Bowman-Birk inhibitor, oleyl disulfide conjugate
37330-34-ODP, Bowman-Birk inhibitor, reaction product with
N-succinimidyl-3-(2-pyridyldithio)propionate and N-Palmityl-2-
pyridyldithiocysteine 171735-25-4DP, reaction products with proteins
254453-83-3P 286365-28-4P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);
BIOL (Biological study); PREP (Preparation); USES (Uses)
(conjugates of hydrophilic mols. with fatty acid or steroid
disulfide derivs. for improving their bioavailabilities)
- IT 285981-92-2P 285981-94-4P
RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological
study); PREP (Preparation); USES (Uses)
(conjugates of hydrophilic mols. with fatty acid or steroid
disulfide derivs. for improving their bioavailabilities)
- IT 57-11-4D, Stearic acid, disulfide derivs., conjugates with
proteins 81-25-4D, disulfide derivs., conjugates with
proteins 83-44-3D, disulfide derivs., conjugates with
proteins 112-80-1D, Oleic acid, disulfide derivs., conjugates
with proteins
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(conjugates of hydrophilic mols. with fatty acid or steroid
disulfide derivs. for improving their bioavailabilities)
- IT 9003-99-ODP, Peroxidase, reaction product with N-succinimidyl-3-(2-
pyridyldithio)propionate and N-Palmityl-2-pyridyldithiocysteine
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);
BIOL (Biological study); PREP (Preparation); USES (Uses)
(horseradish; conjugates of hydrophilic mols. with fatty acid
or steroid disulfide derivs. for improving their bioavailabilities)
- IT 9003-99-0, Peroxidase
RL: RCT (Reactant); RACT (Reactant or reagent)
(horseradish; preparation of conjugates of hydrophilic mols. with
fatty acid or steroid disulfide derivs. for improving their
bioavailabilities)
- IT 52-90-4, L-Cysteine, reactions 83-44-3 1200-22-2, Lipoic acid
2127-03-9, 2,2'-Dithiobis(pyridine) 6066-82-6, N-Hydroxysuccinimide
14464-31-4 16679-58-6, Desmopressin 37330-34-0, Bowman-Birk inhibitor
47931-85-1, Salmon calcitonin 59277-89-3, Acyclovir 68181-17-9, SPDP
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of conjugates of hydrophilic mols. with fatty acid or
steroid disulfide derivs. for improving their bioavailabilities)
- IT 25596-79-6P, Calcitonin (salmon reduced) 37330-34-ODP, Bowman-Birk
inhibitor, reaction product with N-succinimidyl-3-(2-
pyridyldithio)propionate 88442-68-6P 119364-41-9P 171735-25-4P
174069-00-2P 177902-84-0P 285981-91-1P 285981-93-3P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(preparation of conjugates of hydrophilic mols. with fatty acid or
steroid disulfide derivs. for improving their bioavailabilities)

RE.CNT 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Anon; J Pharm Sci 1990, V79(7), P595
- (2) Broadwell, R; Proc Natl Acad Sci USA 1988, V85(2), P632 HCAPLUS
- (3) Brocklehurst, K; Biochem J 1973, V135(1), P573
- (4) Chekhonin, V; FEBS Letters 1991, V287(1,2), P149

Search done by Noble Jarrell

- (5) Chu, Y; Biochem 1994, V33(44), P13087 HCAPLUS
- (6) Conradi, R; Pharm Res 1991, V8(12), P1453 HCAPLUS
- (7) Edwards, P; British Med Bulletin 1978, V34(1), P55 HCAPLUS
- (8) Ekrami, H; A Dissertation, UMI Dissertation Services, Chapter 11 1995, P98
- (9) Ekrami, H; FEBS Letters 1995, V371(3), P283 HCAPLUS
- (10) Eriksson, S; Biochim Biophys Acta E 1970, V212(3), P518 HCAPLUS
- (11) Eriksson, S; Chem Abstracts 1970, P25 HCAPLUS
- (12) Fix, J; Amer J Physiology 1986, V251(3, Prt 1), PG332
- (13) Friden, P; Frontiers In Cerebral Vascular Biology, Transport and Its Regulation 1993, P129 HCAPLUS
- (14) Gonzalez-Mariscal, L; J Membrane Biol 1985, V86(2), P113 HCAPLUS
- (15) Gordon, G; Proc Natl Acad Sci USA 1985, V82(21), P7419 HCAPLUS
- (16) Hashimoto, M; Pharm Res 1989, V6(2), P171 HCAPLUS
- (17) Huang, W; Mol Immunol 1994, V31(15), P1191 HCAPLUS
- (18) Hughes, R; J Pharm Sci 1991, V80(12), P1103 HCAPLUS
- (19) Hughes, R; J Pharm Sci 1992, V81(8), P845 HCAPLUS
- (20) Inagaki, M; Rhinology 1985, V23(3), P213 MEDLINE
- (21) Kabanov, A; Protein Engineering 1989, V3(1), P39 HCAPLUS
- (22) Kajii, H; Life Science 1985, V37(6), P523 HCAPLUS
- (23) Kauvar; US 5599903 1997 HCAPLUS
- (24) Kauvar; US 5679643 1997 HCAPLUS
- (25) Kauvar; US 5763570 1998 HCAPLUS
- (26) Kidron, M; Life Sci 1982, V31(25), P2837 HCAPLUS
- (27) Landolph, J; Transformation Assay of Established Cell Lines: Mechanisms and Application 1985, V67, P185 HCAPLUS
- (28) Lee, V; Crit Rev Ther Drug Carrier Syst 1988, V5(2), P69 HCAPLUS
- (29) Lee, V; Crit Rev Ther Drug Carrier Syst 1991, V8(2), P91 HCAPLUS
- (30) Leone-Bay; US 5629020 1997 HCAPLUS
- (31) Letsinger, R; Proc Natl Acad Sci 1989, V80(17), P6553
- (32) Lowry, O; J Biol Chem 1951, V193, P265 HCAPLUS
- (33) Martins, M; Biochimie 1990, V72(9), P671 HCAPLUS
- (34) McConahey, P; Methods Enzymol 1980, V70(A), P210 MEDLINE
- (35) Mostov, K; Cell 1985, V43(2), P389 HCAPLUS
- (36) Muller, C; International J Pharmaceutics 1989, V57, P41
- (37) Muranishi, S; Pharm Res 1991, V8(5), P649 HCAPLUS
- (38) Naftilan; US 5635380 1997 HCAPLUS
- (39) Reznikoff, C; Cancer Res 1973, V33(12), P3231 MEDLINE
- (40) Reznikoff, C; Cancer Res 1973, V33(12), P3239 HCAPLUS
- (41) Sett, R; J Infectious Dis 1993, V168(4), P994 HCAPLUS
- (42) Shen, W; Adv Drug Delivery Rev 1992, V8(1), P93 HCAPLUS
- (43) Shen, W; Proc Natl Acad Sci USA 1981, V78(12), P7589 HCAPLUS
- (44) Smith, P; Adv Drug Delivery Rev 1992, V8(2,3), P253
- (45) Stn Information Service; File Registry
- (46) Takaori, K; Biochem Biophys Res Commun 1986, V137(2), P682 HCAPLUS
- (47) Taub, M; J Cell Phys 1992, V150(2), P283 HCAPLUS
- (48) Toth, I; J Drug Targeting 1994, V2(3), P217 HCAPLUS
- (49) Ubuka, T; "Synthesis of disulfides related to glutathione and their detection in tissue," Abstract No 156843v 1987 HCAPLUS
- (50) Ubuka, T; (Ganryu Aminosan) Sulfur Amino Acids 1985, V8(1), P153 HCAPLUS
- (51) Uchimi, I; "Glutathione derivatives," Abstract No 54179m 1971 HCAPLUS
- (52) Uchimi, I; "Glutathione derivatives," Abstract No 56440t 1970 HCAPLUS
- (53) Vetvicka, V; Crit Rev Ther Drug Carrier Syst 1988, V5(3), P141 HCAPLUS
- (54) Vitetta, E; J Clin Immunol 1990, V10(6 Suppl), P15S MEDLINE
- (55) Wan, J; J Biol Chem 1992, V267(19), P13446 HCAPLUS
- (56) Wan, J; J Cell Phys 1990, V145(1), P9 HCAPLUS
- (57) Wan, J; Pharm Res 1991, V8(10 Suppl), PS
- (58) Willner; US 5708146 1998 HCAPLUS
- (59) Yavelow, J; Cancer Res 1983, V43(5 Suppl), P2454s MEDLINE
- (60) Yodoya, E; J Pharm Ex Ther 1994, V271(3), P1509 HCAPLUS
- (61) Yoshikawa, H; Pharm Res 1985, V5, P249

IT 285981-92-2P

RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(conjugates of hydrophilic mols. with fatty acid or steroid disulfide derivs. for improving their bioavailabilities)

RN 285981-92-2 HCAPLUS


CN Glycinamide, N-(3-mercapto-1-oxopropyl)-L-tyrosyl-L-phenylalanyl-L-glutamyl-L-asparaginyl-L-cysteinyl-L-prolyl-D-arginyl-, bis(disulfide) with N-[(3 α ,5 β ,12 α)-3,12-dihydroxy-24-oxocholan-24-yl]-L-cysteine (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A




PAGE 1-B



PAGE 1-C




PAGE 2-A




IT 81-25-4D, disulfide derivs., conjugates with proteins
 83-44-3D, disulfide derivs., conjugates with proteins
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (conjugates of hydrophilic mols. with fatty acid or steroid
 disulfide derivs. for improving their bioavailabilities)
 RN 81-25-4 HCAPLUS
 CN Cholan-24-oic acid, 3,7,12-trihydroxy-, (3 α ,5 β ,7 α ,12 α .alph
 a.)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 83-44-3 HCAPLUS
 CN Cholan-24-oic acid, 3,12-dihydroxy-, (3 α ,5 β ,12 α)- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.




IT 83-44-3
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of conjugates of hydrophilic mols. with fatty acid or
 steroid disulfide derivs. for improving their bioavailabilities)
 RN 83-44-3 HCAPLUS
 CN Cholan-24-oic acid, 3,12-dihydroxy-, (3 α ,5 β ,12 α)- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.



IT 174069-00-2P 285981-91-1P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation of conjugates of hydrophilic mols. with fatty acid or
 steroid disulfide derivs. for improving their bioavailabilities)
 RN 174069-00-2 HCAPLUS
 CN 2,5-Pyrrolidinedione, 1-[[(3 α ,5 β ,12 α)-3,12-dihydroxy-24-
 oxocholan-24-yl]oxy]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 285981-91-1 HCAPLUS

CN L-Alanine, N-[(3 α ,5 β ,12 α)-3,12-dihydroxy-24-oxocholan-24-yl]-3-(2-pyridinyldithio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 3 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2000:10612 HCAPLUS

DN 132:73648

ED Entered STN: 06 Jan 2000

TI Lipophilic insulin derivatives soluble at physiological pH with prolonged serum half-lives and biological activity

IN Havelund, Svend; Halstrom, John; Jonassen, Ib; Andersen, Asser Sloth; Markussen, Jan

PA Novo Nordisk A/S, Den.

SO U.S., 47 pp., Cont.-in-part of U.S. 5,750,497.

CODEN: USXXAM

DT Patent

LA English

IC C07K014-62; A61K038-28

INCL 514003000

CC 1-10 (Pharmacology)

Section cross-reference(s): 2

FAN.CNT 3

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

Search done by Noble Jarrell

PI	US 6011007	A	20000104	US 1997-975365	19971120 <--
	ZA 9407187	A	19950317	ZA 1994-7187	19940916 <--
	JP 2000060556	A2	20000229	JP 1999-221632	19940916 <--
	EP 1132404	A2	20010912	EP 2001-112992	19940916 <--
	EP 1132404	A3	20020327		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, LT				
	JP 2002308899	A2	20021023	JP 2001-385921	19940916 <--
	US 5750497	A	19980512	US 1995-400256	19950308 <--
	US 6869930	B1	20050322	US 1999-398365	19990917 <--
	AU 745983	B2	20020411	AU 2000-51960	20000811 <--
	US 2004110664	A1	20040610	US 2002-101454	20020312 <--
PRAI	DK 1993-1044	A	19930917	<--	
	US 1995-400256	A2	19950308	<--	
	US 1994-190829	A	19940202	<--	
	EP 1994-926816	A3	19940916	<--	
	JP 1995-508923	A3	19940916	<--	
	JP 1999-221632	A3	19940916	<--	
	US 1997-975365	A3	19971120	<--	
	US 1999-398365	A1	19990917	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES	
US 6011007	IC	C07K014-62IC	A61K038-28
	INCL	514003000	
US 6011007	NCL	514/003.000; 514/866.000; 530/303.000; 530/304.000	
	ECLA	C07K014/62	<--
EP 1132404	ECLA	C07K014/62	<--
US 5750497	NCL	514/003.000; 514/866.000; 530/304.000	
	ECLA	C07K014/62	<--
US 6869930	NCL	514/003.000; 514/866.000; 530/304.000	
	ECLA	C07K014/62	<--
US 2004110664	NCL	514/003.000	
	ECLA	C07K014/62	<--

OS MARPAT 132:73648

AB Human insulin derivs. with improved solubility at physiol. pH and that retain biol. activity for longer than wild-type human insulin are described. The insulins are substituted at positions A21 and B3 with either being any amino acid except lysine, arginine, or cysteine. The phenylalanine at B1 may be deleted and the amino acid at position B30 may be deleted or substituted by any amino acid except lysine, arginine, or cysteine or by another amino acid that is lipophilic having a C10-24 side chain. If B30 is deleted or substituted, lysineB29 is modified by a carboxylic acid connected to the ε-amino group. When B30 is threonine or alanine and A21 and B3 are both asparagine, and phenylalanineB1 is present, then the insulin derivative is always present as a Zn₂ complex.

ST human insulin sequence acylation diabetes pharmaceutical; lipophilic insulin deriv antidiabetic

IT Carboxylic acids, reactions

RL: RCT (Reactant); RACT (Reactant or reagent)

(C5, insulins modified with; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Solubility

(at physiol. pH; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Carboxylic acids, reactions

RL: RCT (Reactant); RACT (Reactant or reagent)

(dicarboxylic, C<6, insulin modification by; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT cDNA sequences

(for insulin analogs of human; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Drug delivery systems

(injections, insulin; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Antidiabetic agents

(insulin analogs as; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Acetyl group
Formyl group
(insulin derivs. containing; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Fatty acids, reactions
RL: RCT (Reactant); RACT (Reactant or reagent)
(insulin derivs. containing; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Protein sequences
(lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Lipophilicity
(of insulin derivs.; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Plasmids
(pAK-series and pKFN1627 and pEA-series; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT Functional groups
(propionyl, insulin derivs. containing; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 14464-31-4 69888-86-4 88404-23-3 104943-24-0 165893-02-7
165893-03-8 168986-19-4 168986-20-7 169142-69-2
RL: RCT (Reactant); RACT (Reactant or reagent)
(acylation of insulin derivs. using; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 169148-55-4DP, zinc complexes 169148-56-5DP, zinc complexes
169148-57-6P 169148-58-7P 169148-59-8P 169148-60-1P 169148-61-2DP,
zinc complexes 169148-62-3DP, zinc complexes 169148-63-4P
169148-64-5P 169148-65-6P 169148-66-7P 169148-67-8P 169148-68-9P
169148-69-0P 169148-70-3P 169148-71-4P 169148-72-5DP, zinc complexes
169148-73-6P 169148-74-7DP, zinc complexes 169148-75-8DP, zinc
complexes 169535-16-4P 169535-18-6P 169535-20-0P 169535-22-2P
169535-24-4P 169535-26-6P 169535-28-8P 169535-30-2P 169535-32-4P
169535-34-6P 169535-36-8P 169535-38-0P
RL: BPN (Biosynthetic preparation); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(amino acid sequence; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 11061-68-0D, Insulin (human), amino acid-substituted, derivatized
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 51-49-0DP, conjugates with insulin 108-30-5DP,
conjugates with insulin 110-15-6DP, Butanedioic acid,
conjugates with insulin, preparation 143-07-7DP, Dodecanoic
acid, conjugates with insulin, preparation 544-63-8DP,
Tetradecanoic acid, conjugates with insulin, preparation
638-53-9DP, Tridecanoic acid, conjugates with insulin
7145-63-3DP, conjugates with insulin 7452-59-7DP,
conjugates with insulin 7769-79-1DP, conjugates with
insulin 14565-47-0DP, conjugates with insulin 17702-88-4DP,
conjugates with insulin 22102-66-5DP, conjugates with
insulin 35237-37-7DP, conjugates with insulin 68528-80-3DP,
conjugates with insulin 104211-94-1DP,
conjugates with insulin 141537-81-7DP, conjugates with
insulin 158627-30-6DP, conjugates with insulin
168986-14-9DP, conjugates with insulin 168986-15-0DP,
conjugates with insulin 168986-16-1DP, conjugates with
insulin
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(lipophilic insulin derivs. soluble at physiol. pH with prolonged serum

half-lives and biol. activity)

IT 23713-49-7DP, Zinc dication, complexes with insulin derivs., biological studies
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 169535-21-1P 169535-23-3P, DNA (Saccharomyces cerevisiae synthetic signal peptide LaC212spx3 fusion protein with synthetic peptide fusion protein with human insulin A chain [21-glycine] fusion protein with human insulin B-chain [3-aspartic acid]-specifying cDNA plus flanks)
 169535-27-7P 169535-29-9P 169535-33-5P 169535-35-7P 169535-37-9P 169535-39-1P
 RL: BPN (Biosynthetic preparation); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (nucleotide sequence; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 169535-17-5, DNA (Saccharomyces cerevisiae clone pAK188 synthetic signal peptide LaC212spx3 fusion protein with human clone pAK188 1-29-insulin B-chain fusion protein with synthetic clone pAK188 5-amino acid peptide fusion protein with human clone pAK188 insulin A-chain-specifying plus flanks) 169535-19-7 169535-25-5, DNA (Saccharomyces cerevisiae synthetic signal peptide LaC212spx3 fusion protein with human insulin A-chain [21-glycine] fusion protein with human insulin B-chain [3-aspartic acid]-specifying cDNA plus flanks)
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (nucleotide sequence; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 24424-99-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (protecting group; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 222586-86-9, 3: PN: US6011007 SEQID: 3 unclaimed DNA 222586-87-0, 4: PN: US6011007 SEQID: 4 unclaimed DNA 222586-88-1, 5: PN: US6011007 SEQID: 5 unclaimed DNA 222586-89-2, 6: PN: US6011007 SEQID: 6 unclaimed DNA 222586-90-5, 7: PN: US6011007 SEQID: 7 unclaimed DNA 222586-91-6, 8: PN: US6011007 SEQID: 8 unclaimed DNA 222586-92-7, 9: PN: US6011007 SEQID: 9 unclaimed DNA 222586-93-8 222586-94-9 222586-95-0 222586-96-1 222586-98-3 222587-00-0 222587-09-9
 RL: PRP (Properties)
 (unclaimed nucleotide sequence; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

IT 253597-47-6 253597-48-7
 RL: PRP (Properties)
 (unclaimed protein sequence; lipophilic insulin derivs. soluble at physiol. pH with prolonged serum half-lives and biol. activity)

RE.CNT 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Anon; GB 1492997 1977 HCAPLUS
- (2) Anon; JP 5767548 1982
- (3) Anon; WO 9112817 1991 HCAPLUS
- (4) Anon; Prescription Products Guide 1992, P942
- (5) Brange And Langkjaer; Chemical Stability of Insulin Acta Pharm Nord 1992, V4(3), P149
- (6) Breddam; US 4645740 1987 HCAPLUS
- (7) Doerge; Wilson and Gisvold's textbook of organic medicinal and pharmaceutical chemistry 1982, P774
- (8) Foye, W; Principles of Medicinal Chemistry 1974, P563
- (9) Gammelhoft; Phys Rev 1984, V64, P1321
- (10) Grant; US 3823125 1974 HCAPLUS
- (11) Haas; US 3528960 1970 HCAPLUS
- (12) Kurtz; Diabetologia 1983, V25, P322 MEDLINE
- (13) Lindsay; US 3950517 1976 HCAPLUS
- (14) Lindsay And Shall; The Acetylation of Insulin Biochem 1991, V121, P737

- (15) Marble, A; Joslin's Diabetes Mellitus, 12th Edition 1985, P380
- (16) Markussen; US 5008241 1991 HCAPLUS
- (17) Markussen; Prot Eng 1987, V1, P205 HCAPLUS
- (18) Markussen; Prot Eng 1988, V2, P157 HCAPLUS
- (19) Mims Annual; Section 6d "Insulin Preparations" 1991
- (20) Mims Annual; Section 6d "Insulin Preparations" 1993
- (21) Panayotis; US 5208217 1993 HCAPLUS
- (22) Samuel; Clin Exp Immunol 1978, V33, P252 HCAPLUS
- (23) Schade; Excerpta Medica 1983, P7
- (24) Schlichtkrull, J; "Insulin Crystals" (Ejnar Munksgaard) 1958, P21
- (25) Smyth; US 3868356 1975 HCAPLUS

IT 168986-19-4


RL: RCT (Reactant); RACT (Reactant or reagent)

(acylation of insulin derivs. using; lipophilic insulin derivs. soluble at
physiol. pH with prolonged serum half-lives and biol. activity)

RN 168986-19-4 HCAPLUS

CN Pentanoic acid, 5-[(2,5-dioxo-1-pyrrolidinyl)oxy]-4-[[(3 α ,5 β)-3-
hydroxy-24-oxocholan-24-yl]amino]-5-oxo-, 1,1-dimethylethyl ester, (4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 104211-94-1DP, conjugates with insulin


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)

(lipophilic insulin derivs. soluble at physiol. pH with prolonged serum
half-lives and biol. activity)

RN 104211-94-1 HCAPLUS

CN 2,5-Pyrrolidinedione, 1-[[(3 α ,5 β)-3-hydroxy-24-oxocholan-24-
yl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 4 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1999:722480 HCAPLUS
 DN 132:313463
 ED Entered STN: 12 Nov 1999
 TI Thermosensitive self-aggregates prepared from cholic acid-
 conjugated amine-terminated poly(N-isopropylacrylamide) for drug
 delivery
 AU Kim, I. S.; Kim, S. H.
 CS College of Pharmacy, Chosun University, Kwangju, 501-759, S. Korea
 SO Proceedings of the International Symposium on Controlled Release of
 Bioactive Materials (1999), 26th, 791-792
 CODEN: PCRMEY; ISSN: 1022-0178
 PB Controlled Release Society, Inc.
 DT Journal
 LA English
 CC 63-5 (Pharmaceuticals)
 AB Polymer micelles composed of cholic acid and amine-terminated
 thermoresponsive poly(N-isopropylacrylamide) were prepared and showed
 reversible thermal transition. Drug delivery systems using these
 thermosensitive micelles can be used for the site-specific drug delivery
 by modulating the temperature at the target site.
 ST polyisopropylacrylamide nanoparticle micelle cholate drug delivery
 IT Drug delivery systems
 (nanoparticles, controlled-release; thermosensitive self-aggregates
 prepared from cholic acid-conjugated amine-terminated
 poly(N-isopropylacrylamide) for drug delivery)
 IT Micelles
 Self-association
 (thermosensitive self-aggregates prepared from cholic acid-
 conjugated amine-terminated poly(N-isopropylacrylamide) for
 drug delivery)
 IT 81-25-4DP, Cholic acid, reaction products with amine-terminated
 poly(isopropylacrylamide) 25189-55-3DP, Poly(N-isopropylacrylamide),
 amine-terminated, reaction products with cholic acid
 RL: PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use);
 BIOL (Biological study); PREP (Preparation); USES (Uses)
 (thermosensitive self-aggregates prepared from cholic acid-
 conjugated amine-terminated poly(N-isopropylacrylamide) for
 drug delivery)
 IT 53-86-1, Indomethacin
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES
 (Uses)
 (thermosensitive self-aggregates prepared from cholic acid-

conjugated amine-terminated poly(N-isopropylacrylamide) for drug delivery)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE


- (1) Anton, P; Makromol Chem 1993, V194, P1 HCAPLUS
- (2) Bae, Y; J Controlled Release 1989, V9, P271 HCAPLUS
- (3) Gao, Z; Macromolecules 1993, V26, P7353 HCAPLUS
- (4) Guenoun, P; Macromolecules 1996, V29, P3965 HCAPLUS
- (5) Xu, R; Macromolecules 1991, V24, P87 HCAPLUS

IT 81-25-4DP, Cholic acid, reaction products with amine-terminated poly(isopropylacrylamide)
RL: PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(thermosensitive self-aggregates prepared from cholic acid-conjugated amine-terminated poly(N-isopropylacrylamide) for drug delivery)

RN 81-25-4 HCAPLUS

CN Cholan-24-oic acid, 3,7,12-trihydroxy-, (3 α ,5 β ,7 α ,12 α .alph a.)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 5 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1999:708452 HCAPLUS

DN 131:314185

ED Entered STN: 05 Nov 1999

TI Active hedgehog protein conjugate, process for its production and use

IN Esswein, Angelika; Lang, Kurt; Rueger, Petra; Seytter, Tilmann

PA Roche Diagnostics G.m.b.H., Germany

SO Eur. Pat. Appl., 25 pp.

CODEN: EPXXDW

DT Patent

LA English

IC ICM C07K014-47

ICA C07K019-00

CC 63-5 (Pharmaceuticals)

FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 953576	A1	19991103	EP 1999-108032	19990423 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	EP 953575	A1	19991103	EP 1998-107911	19980430 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	NZ 335385	A	20000929	NZ 1999-335385	19990426 <--
	MX 9903976	A	20000630	MX 1999-3976	19990428 <--
	SG 80028	A1	20010417	SG 1999-2117	19990428 <--
	US 6468978	B1	20021022	US 1999-301199	19990428 <--
	CA 2269221	AA	19991030	CA 1999-2269221	19990429 <--

Search done by Noble Jarrell

NO 9902090	A	19991101	NO 1999-2090	19990429 <--
ZA 9903009	A	19991101	ZA 1999-3009	19990429 <--
CN 1233616	A	19991103	CN 1999-106302	19990429 <--
AU 9925009	A1	19991111	AU 1999-25009	19990429 <--
AU 719797	B2	20000518		
JP 2000053699	A2	20000222	JP 1999-125005	19990430 <--
JP 3433136	B2	20030804		
BR 9903169	A	20001017	BR 1999-3169	19990430 <--
US 2003139574	A1	20030724	US 2002-278060	20021021 <--
US 6818623	B2	20041116		
PRAI EP 1998-107911	A	19980430	<--	
EP 1998-116733	A	19980903	<--	
US 1999-301199	A1	19990428	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES	
EP 953576	ICM	C07K014-47	
	ICA	C07K019-00	
EP 953576	ECLA	C07K014/47	<--
EP 953575	ECLA	C07K014/47	<--
US 6468978	NCL	514/021.000; 514/012.000; 530/350.000; 530/408.000; 530/409.000; 530/410.000	
	ECLA	C07K014/47	<--
US 2003139574	NCL	514/021.000; 514/012.000; 530/350.000; 530/408.000; 530/409.000; 530/410.000	
	ECLA	C07K014/47	<--

AB A hedgehog conjugate is disclosed which is characterized in that it contains: (a) a polypeptide composed of 10 to 30 hydrophobic amino acids and/or amino acids which form transmembrane helixes and are pos. charged, (b) 1 to 4 aliphatic, saturated or unsatd. hydrocarbon residues with a chain length of 10 to 24 C atoms and with a hydrophobic action or (c) a hydrophobic thio compound covalently bound to a hedgehog protein and which has a several-fold increased activity and is suitable as a pharmaceutical agent.

ST hedgehog protein lipid conjugate drug

IT Polysaccharides, uses

RL: NUU (Other use, unclassified); USES (Uses)
(acidic; active hedgehog protein conjugates for therapeutic use)

IT DNA sequences

Detergents

Drug delivery systems

Molecular cloning

Stabilizing agents

(active hedgehog protein conjugates for therapeutic use)

IT Primers (nucleic acid)

RL: PRP (Properties)

(active hedgehog protein conjugates for therapeutic use)

IT Alcohols, biological studies

RL: BAC (Biological activity or effector, except adverse); BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(alkyl, hedgehog protein conjugates; active hedgehog protein conjugates for therapeutic use)

IT Fatty acids, biological studies

RL: BAC (Biological activity or effector, except adverse); BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(hedgehog protein conjugates; active hedgehog protein conjugates for therapeutic use)

IT Hedgehog protein

RL: BAC (Biological activity or effector, except adverse); BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP

(Preparation); USES (Uses)
 (lipid conjugates; active hedgehog protein conjugates
 for therapeutic use)

IT Dimerization
 (of human sonic hedgehog protein; active hedgehog protein
 conjugates for therapeutic use)

IT Hedgehog protein
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP
 (Preparation)
 (sonic, cloning of human; active hedgehog protein conjugates
 for therapeutic use)

IT 57-10-3DP, Palmitic acid, hedgehog protein conjugates
 57-11-4DP, Stearic acid, hedgehog protein conjugates
 60-33-3DP, Linoleic acid, hedgehog protein conjugates
 112-80-1DP, Oleic acid, hedgehog protein conjugates
 112-85-6DP, Behenic acid, hedgehog protein conjugates
 143-07-7DP, Lauric acid, hedgehog protein conjugates
 373-49-9DP, Palmitoleic acid, hedgehog protein conjugates
 463-40-1DP, Linolenic acid, hedgehog protein conjugates
 506-30-9DP, Arachidic acid, hedgehog protein conjugates
 506-32-1DP, Arachidonic acid, hedgehog protein conjugates
 544-63-8DP, Myristic acid, hedgehog protein conjugates
 1249-81-6DP, Thiocholesterol, hedgehog protein conjugates
 RL: BAC (Biological activity or effector, except adverse); BPN
 (Biosynthetic preparation); BSU (Biological study, unclassified); PRP
 (Properties); THU (Therapeutic use); BIOL (Biological study); PREP
 (Preparation); USES (Uses)
 (active hedgehog protein conjugates for therapeutic use)

IT 145-63-1, Suramin 9005-49-6, Heparin, uses
 RL: NUU (Other use, unclassified); USES (Uses)
 (active hedgehog protein conjugates for therapeutic use)

IT 361-09-1, Sodium cholate 2281-11-0, Zwittergent 3-16
 9002-93-1, Triton x 100 9005-65-6, Tween 80 14933-09-6, Zwittergent
 3-14 41444-50-2, Octyl glucoside 75621-03-3, Chaps
 RL: NUU (Other use, unclassified); PRP (Properties); USES (Uses)
 (active hedgehog protein conjugates for therapeutic use)

IT 3867-67-2P 17450-31-6P 26227-65-6P 60988-34-3P 69205-88-5P
 69205-89-6P 136911-91-6P 247900-73-8P 247900-74-9P
 247900-75-0P 247900-76-1P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (active hedgehog protein conjugates for therapeutic use)

IT 1763-10-6, Palmitoyl-CoA
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (coupling agent; active hedgehog protein conjugates for
 therapeutic use)


RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Beachy, P; WO 9830576 A 1998 HCAPLUS
 (2) Farese, R; TRENDS IN GENETICS 1998, V14(3), P115 HCAPLUS
 (3) Hammerschmidt, M; TRENDS IN GENETICS 1997, V13(1), P14 HCAPLUS
 (4) Hancock; CELL 1990, V63, P133 HCAPLUS
 (5) Harvard College; WO 9518856 A 1995 HCAPLUS
 (6) Mohler; DEVELOPMENT 1992, V115, P957 HCAPLUS
 (7) Porter; CELL 1996, V86, P21 HCAPLUS
 (8) Porter; SCIENCE 1996, V274, P255 HCAPLUS

IT 361-09-1, Sodium cholate
 RL: NUU (Other use, unclassified); PRP (Properties); USES (Uses)
 (active hedgehog protein conjugates for therapeutic use)

RN 361-09-1 HCAPLUS

CN Cholan-24-oic acid, 3,7,12-trihydroxy-, monosodium salt,
 (3 α ,5 β ,7 α ,12 α)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● Na

IT 247900-74-9P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (active hedgehog protein conjugates for therapeutic use)
 RN 247900-74-9 HCAPLUS
 CN 1H-Imidazole, 1-(24-oxocholan-24-yl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 6 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1998:208387 HCAPLUS
 DN 128:286354
 ED Entered STN: 13 Apr 1998
 TI Methods and compositions for lipidization of hydrophilic molecules
 IN Shen, Wei-Chiang; Wang, Jinghua
 PA University of Southern California, USA; Shen, Wei-Chiang; Wang, Jinghua
 SO PCT Int. Appl., 52 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM A61K
 CC 63-6 (Pharmaceuticals)
 Section cross-reference(s): 1

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9813007	A2	19980402	WO 1997-US17282	19970926 <--
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG,				

Search done by Noble Jarrell

UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR,
 GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA,
 GN, ML, MR, NE, SN, TD, TG

CA 2267179	AA	19980402	CA 1997-2267179	19970926 <--
AU 9745967	A1	19980417	AU 1997-45967	19970926 <--
AU 737865	B2	20010906		
CN 1235594	A	19991117	CN 1997-199191	19970926 <--
CN 1127477	B	20031112		
EP 1023316	A2	20000802	EP 1997-944483	19970926 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
BR 9712128	A	20001212	BR 1997-12128	19970926 <--
JP 2002515883	T2	20020528	JP 1998-515933	19970926 <--
NO 9901465	A	19990510	NO 1999-1465	19990325 <--
KR 2000048608	A	20000725	KR 1999-702543	19990325 <--
PRAI US 1996-721306	A	19960926	<--	
US 1997-49499P	P	19970613	<--	
US 1996-77177P	P	19960926	<--	
WO 1997-US17282	W	19970926	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES	
WO 9813007	ICM	A61K	
WO 9813007	ECLA	A61K047/48H4	<--
OS	MARPAT 128:286354		
AB	Fatty acid derivs. of disulfide-containing compds. (for example, disulfide-containing peptides or proteins) comprising fatty acid-conjugated products with a disulfide linkage are employed for delivery of the compds. to mammalian cells. This modification markedly increases the absorption of the compds. by mammalian cells relative to the rate of absorption of the unconjugated compds., as well as prolonging blood and tissue retention of the compds. Moreover, the disulfide linkage in the conjugate is quite labile in vivo and thus facilitates intracellular or extracellular release of the intact compds. from the fatty acid moieties. N-palmitoyl-2-pyridyldithiocysteine was prepared and conjugated to BBI hydrophilic protein and its transport and biodistribution studied.		
ST	lipidization hydrophilic compd delivery; fatty acid deriv protein peptide delivery		
IT	Drug delivery systems (lipidization of hydrophilic mols. for peptide or protein delivery)		
IT	Disulfides RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (lipidization of hydrophilic mols. for peptide or protein delivery)		
IT	Drug delivery systems (liposomes; lipidization of hydrophilic mols. for peptide or protein delivery)		
IT	Proteins, general, biological studies RL: BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses) (reaction products with palmitoylpyridyldithiocysteine, lipidization of hydrophilic mols. for peptide or protein delivery)		
IT	Oligonucleotides Peptides, biological studies Proteins, specific or class RL: BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses) (reaction products, with fatty acids; lipidization of hydrophilic mols. for peptide or protein delivery)		
IT	Fatty acids, biological studies RL: BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)		

(reaction products, with proteins and peptides; lipidization of hydrophilic mols. for peptide or protein delivery)

IT 112-80-1DP, Oleic acid, derivative, reaction products with peptides and proteins 1200-22-2DP, Lipoic acid, reaction products with acyclovir and palmitic acid derivative 9003-99-0DP, Peroxidase, reaction products with palmitic acid derivative 16679-58-6DP, Desmopressin, reaction products with palmitic acid derivative 47931-85-1DP, Salmon calcitonin, reaction products with palmitic acid derivative 59277-89-3DP, Acyclovir, reaction products with lipoic acid and palmitic acid derivative 171735-25-4DP, reaction products with peptides and proteins 174069-00-2DP, reaction products with palmitic acid derivative

RL: BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)

(lipidization of hydrophilic mols. for peptide or protein delivery)

IT 52-90-4, L-Cysteine, reactions 83-44-3, Deoxycholic acid 2127-03-9, Pyridine, 2,2'-Dithiobis- 6066-82-6, N-Hydroxysuccinimide 68181-17-9, SPDP

RL: RCT (Reactant); RACT (Reactant or reagent)

(lipidization of hydrophilic mols. for peptide or protein delivery)

IT 14464-31-4P, N-Hydroxysuccinimide palmitate 88442-68-6P 171735-25-4P 174069-00-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(lipidization of hydrophilic mols. for peptide or protein delivery)


IT 174069-00-2DP, reaction products with palmitic acid derivative
RL: BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)

(lipidization of hydrophilic mols. for peptide or protein delivery)

RN 174069-00-2 HCAPLUS

CN 2,5-Pyrrolidinedione, 1-[[[(3 α ,5 β ,12 α)-3,12-dihydroxy-24-oxocholan-24-yl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 83-44-3, Deoxycholic acid


RL: RCT (Reactant); RACT (Reactant or reagent)

(lipidization of hydrophilic mols. for peptide or protein delivery)

RN 83-44-3 HCAPLUS


CN Cholan-24-oic acid, 3,12-dihydroxy-, (3 α ,5 β ,12 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 174069-00-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (lipidization of hydrophilic mols. for peptide or protein delivery)
 RN 174069-00-2 HCAPLUS
 CN 2,5-Pyrrolidinedione, 1-[[[(3 α ,5 β ,12 α)-3,12-dihydroxy-24-oxocholan-24-yl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 7 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1997:463447 HCAPLUS
 DN 127:99659
 ED Entered STN: 24 Jul 1997
 TI Oral peptide delivery using the intestinal bile acid transporter
 AU Swaan, P. W.; Szoka, F. C., Jr.; Oie, S.
 CS University of California at San Francisco, CA, 94143-0446, USA
 SO Proceedings of the International Symposium on Controlled Release of
 Bioactive Materials (1997), 24th, 7-8
 CODEN: PCRMEY; ISSN: 1022-0178
 PB Controlled Release Society, Inc.
 DT Journal
 LA English
 CC 63-5 (Pharmaceuticals)
 Section cross-reference(s): 1
 AB The intestinal absorption of peptides was increased by coupling to the 24
 position of the steroid nucleus in cholic acid.
 ST bile acid peptide delivery intestine
 IT Drug delivery systems
 Intestine

Search done by Noble Jarrell

(oral peptide delivery using intestinal bile acid transporter)

IT Bile acids
Peptides, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(oral peptide delivery using intestinal bile acid transporter)


IT 81-25-4D, Cholic acid, peptide conjugates
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(oral peptide delivery using intestinal bile acid transporter)

IT 81-25-4D, Cholic acid, peptide conjugates
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(oral peptide delivery using intestinal bile acid transporter)

RN 81-25-4 HCAPLUS

CN Cholan-24-oic acid, 3,7,12-trihydroxy-, (3 α ,5 β ,7 α ,12 α .alph
a.)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 8 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
AN 1997:443336 HCAPLUS
DN 127:55909
ED Entered STN: 17 Jul 1997
TI Sulfate conjugates of ursodeoxycholic acid, and their beneficial
use in inflammatory disorders and other applications
IN Setchell, Kenneth D. R.
PA Children's Hospital Medical Center, Philadelphia, USA
SO PCT Int. Appl., 68 pp.
CODEN: PIXXD2
DT Patent
LA English
IC ICM A61K031-575
CC 63-6 (Pharmaceuticals)
Section cross-reference(s): 1

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9718816	A2	19970529	WO 1996-US18487	19961119 <--
	WO 9718816	A3	19970626		
	W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
	US 5763435	A	19980609	US 1995-560992	19951121 <--
	CA 2238040	AA	19970529	CA 1996-2238040	19961119 <--
	CA 2238040	C	20040713		
	AU 9677377	A1	19970611	AU 1996-77377	19961119 <--
	AU 709594	B2	19990902		
	EP 871452	A2	19981021	EP 1996-940516	19961119 <--

Search done by Noble Jarrell

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, FI

CN 1211185	A	19990317	CN 1996-199708	19961119 <--
JP 2000500764	T2	20000125	JP 1997-519828	19961119 <--
BR 9611606	A	20001024	BR 1996-11606	19961119 <--
PL 186393	B1	20040130	PL 1996-326931	19961119 <--
US 6251884	B1	20010626	US 1998-28036	19980224 <--
NO 9802281	A	19980708	NO 1998-2281	19980519 <--
PRAI US 1995-560992	A	19951121	<--	
WO 1996-US18487	W	19961119	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES	
WO 9718816	ICM	A61K031-575	
WO 9718816	ECLA	A61K031/575	<--
US 5763435	NCL	514/182.000	
	ECLA	A61K031/575	<--
US 6251884	NCL	514/182.000; 514/169.000	
	ECLA	A61K031/575	<--
AB	Pharmaceutically acceptable compns. including a sulfate of ursodeoxycholic acid (I), glyoursodeoxycholic acid, or tauroursodeoxycholic acid and a pharmacol. acceptable carrier are useful for treatment of mammals for disorders including inflammation of the gastrointestinal tract, colon cancer, rectum cancer, ulcerative colitis, adenomatous polyps, familial polyposis, hepatitis, etc. These compns. may be used to improve liver function or serum biochem. in liver disease, to increase bile flow, or to decrease biliary secretion of phospholipid or cholesterol. An isolated organ may be maintained in vitro by perfusion with a I sulfate. Thus, I was condensed with tert-butyldimethylsilyl chloride to form the 3-tert-butyldimethylsilyl ether, then with Ac2O to form I 3-tert-butyldimethylsilyl ether 7-acetate, hydrolyzed with HCl to I 7-acetate, condensed with ClSO3H to form I 7-acetate 3-sulfate, converted to the di-Na salt, and saponified with methanolic NaOH to I 3-sulfate.		
ST	ursodeoxycholate sulfate inflammation inhibitor; colon cancer ursodeoxycholate sulfate; rectum cancer ursodeoxycholate sulfate; liver disease ursodeoxycholate sulfate		
IT	Intestine, neoplasm (colon, inhibitors; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)		
IT	Antitumor agents (colon; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)		
IT	Digestive tract (disease, inflammation; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)		
IT	Anti-inflammatory agents (gastrointestinal; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)		
IT	Solutions (isotonic solns., for organ perfusion; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)		
IT	Intestine (large, disease, inflammation; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)		
IT	Bile acids RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process) (metabolism of; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)		
IT	Intestine Kidney Lung Organ preservation (perfusion fluid for; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)		
IT	Intestine, neoplasm		

- (polyp, adenomatous; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT Intestine, neoplasm
Intestine, neoplasm
(rectum, inhibitors; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT Antitumor agents
Antitumor agents
(rectum; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT Phospholipids, biological studies
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(secretion of, in bile; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT Intestine, disease
Intestine, disease
(small, inflammation; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT Cholagogues
Hepatitis
Liver, disease
(sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT Intestine, disease
(ulcerative colitis; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT Biological transport
(uptake, of ursodeoxycholic acid by intestine, inhibition of; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT 81-25-4, Cholic acid 83-44-3, Deoxycholic acid 434-13-9, Lithocholic acid
RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
(formation from ursodeoxycholate; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT 128-13-2, Ursodeoxycholic acid
RL: BPR (Biological process); BSU (Biological study, unclassified); RCT (Reactant); BIOL (Biological study); PROC (Process); RACT (Reactant or reagent)
(intestinal absorption of; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT 57-88-5, Cholesterol, biological studies
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(secretion of, in bile; sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT 71781-68-5P 191286-16-5P 191286-18-7P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT 68780-73-4, Ursodeoxycholic acid 3-sulfate 74723-13-0, Tauroursodeoxycholic acid 3-sulfate 74723-14-1 74723-15-2 74723-16-3 88426-32-8 109333-29-1 133429-88-6, Glycoursodeoxycholic acid 3-sulfate 191286-12-1
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)
- IT 474-25-9, Chenodeoxycholic acid 2393-58-0, α -Muricholic acid 2393-59-1, β -Muricholic acid 6830-03-1, ω -Muricholic acid 114183-56-1 114183-57-2 163750-00-3

RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
 (sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)

IT 18162-48-6, tert-Butyldimethylsilyl chloride
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)


IT 71781-57-2P 71781-58-3P 75672-25-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)

IT 163750-00-3
 RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
 (sulfate conjugates of ursodeoxycholic acid for treatment of inflammatory disorders)

RN 163750-00-3 HCAPLUS

CN Chol-22-en-24-oic acid, 3,7-dihydroxy-, (3 α ,5 β ,7 β)- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry unknown.



L40 ANSWER 9 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1997:372273 HCAPLUS
 DN 126:347323
 ED Entered STN: 14 Jun 1997
 TI Buccal delivery of glucagon-like insulinotropic peptides (GLPs)
 IN Heiber, Sonia J.; Ebert, Charles D.; Gutniak, Mark K.
 PA Theratech, Inc., USA
 SO PCT Int. Appl., 55 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM A61K009-70
 ICS A61L015-16
 CC 63-6 (Pharmaceuticals)

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9715296	A1	19970501	WO 1996-US16890	19961022 <--
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI				
US 5766620	A	19980616	US 1995-553807	19951023 <--

Search done by Noble Jarrell

CA 2235369	AA	19970501	CA 1996-2235369	19961022 <--
AU 9674647	A1	19970515	AU 1996-74647	19961022 <--
AU 716038	B2	20000217		
EP 859606	A1	19980826	EP 1996-936815	19961022 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
CN 1202820	A	19981223	CN 1996-198618	19961022 <--
BR 9611139	A	19990406	BR 1996-11139	19961022 <--
JP 11513982	T2	19991130	JP 1996-516712	19961022 <--
TW 416854	B	20010101	TW 1996-85112962	19961022 <--
ZA 9608909	A	19970528	ZA 1996-8909	19961023 <--
US 5863555	A	19990126	US 1997-964731	19971105 <--
PRAI US 1995-553807	A	19951023	<--	
WO 1996-US16890	W	19961022	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES	
WO 9715296	ICM	A61K009-70	
	ICS	A61L015-16	
US 5766620	NCL	424/436.000; 424/435.000; 514/772.300; 514/772.600; 514/774.000; 514/777.000; 514/781.000	
	ECLA	A61K009/00M18D; A61K038/26	<--
US 5863555	NCL	424/435.000; 514/772.300; 514/772.600; 514/774.000; 514/777.000; 514/781.000	
	ECLA	A61K038/26	<--

AB Drug delivery systems for administering a GLP to the buccal mucosa for transmucosal drug delivery comprise a drug composition containing effective amts. of the GLP and a permeation enhancer, and means for maintaining the drug composition in a drug-transferring relation with the buccal mucosa. These systems can be in free form, such as creams, gels, and ointments, or can comprise a device of determined phys. form, such as tablets, patches, and troches. A preferred GLP is GLP-1(7-36) amide. Thus, a gingival bilayer tablet was prepared comprising an active layer and an adhesive layer. The adhesive layer was prepared by mixing polyethylene oxide 70, Carbopol 934P 20, and compressible xylitol/CM-cellulose filler 10 weight parts, granulating with EtOH, sieving, drying, mixing with stearic acid 0.25 and mint flavor 0.06 weight%, and compression. To prepare the active layer, mannitol 49.39, hydroxypropylcellulose 34.33, and Na taurocholate 15.00 weight% were mixed, granulated with EtOH, sieved, dried, combined with GLP-1(7-36) amide 0.91, FD&C Yellow Number 6HT 0.06, Mg stearate 0.25, and mint flavor 0.06 weight%; 50 mg of this mixture was compressed onto 50 mg adhesive layer.

ST glucagonlike insulintropic peptide buccal tablet; mouth absorption
glucagonlike insulintropic peptide

IT Alcohols, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(C2-3, permeation enhancers; buccal delivery of glucagon-like insulintropic peptides)

IT Glycols, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(C3-4; buccal delivery of glucagon-like insulintropic peptides)

IT Polyoxyalkylenes, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(adhesive containing; buccal delivery of glucagon-like insulintropic peptides)

IT Caseins, biological studies
Gelatins, biological studies
Polyethers, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(adhesives containing; buccal delivery of glucagon-like insulintropic peptides)

IT Adhesives
(biol.; buccal delivery of glucagon-like insulintropic peptides)

IT Antidiabetic agents
Gingiva
Permeation enhancers
(buccal delivery of glucagon-like insulintropic peptides)

- IT Sulfonylureas
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(buccal delivery of glucagon-like insulinotropic peptides)
- IT Drug delivery systems
(buccal; buccal delivery of glucagon-like insulinotropic peptides)
- IT Steroids, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(detergents, as permeation enhancers; buccal delivery of glucagon-like insulinotropic peptides)
- IT Cell membrane
(disrupting agents for; buccal delivery of glucagon-like insulinotropic peptides)
- IT Drug delivery systems
(gels; buccal delivery of glucagon-like insulinotropic peptides)
- IT Polymers, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(hydrophilic, adhesive containing; buccal delivery of glucagon-like insulinotropic peptides)
- IT Drug delivery systems
(lozenges; buccal delivery of glucagon-like insulinotropic peptides)
- IT Mouth
(mucosa; buccal delivery of glucagon-like insulinotropic peptides)
- IT Drug delivery systems
(ointments, creams; buccal delivery of glucagon-like insulinotropic peptides)
- IT Drug delivery systems
(ointments; buccal delivery of glucagon-like insulinotropic peptides)
- IT Chelating agents
Solvents
Surfactants
(permeation enhancers; buccal delivery of glucagon-like insulinotropic peptides)
- IT Bile salts
Fatty acids, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(permeation enhancers; buccal delivery of glucagon-like insulinotropic peptides)
- IT Vinyl compounds, biological studies
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(polymers, adhesives containing; buccal delivery of glucagon-like insulinotropic peptides)
- IT Detergents
(steroidal, as permeation enhancers; buccal delivery of glucagon-like insulinotropic peptides)
- IT Drug delivery systems
Drug delivery systems
(tablets, buccal; buccal delivery of glucagon-like insulinotropic peptides)
- IT 79-10-7D, 2-Propenoic acid, esters, polymers, biological studies
79-10-7D, 2-Propenoic acid, polymers, biological studies 557-75-5D,
Ethenol, polymers, biological studies 9000-30-0, Guar gum 9000-69-5,
Pectin 9003-39-8, PVP 9004-32-4 9004-54-0, Dextran, biological
studies 9004-57-3, Ethylcellulose 9004-62-0, Hydroxyethylcellulose
9004-64-2, Hydroxypropylcellulose 9004-65-3,
Hydroxypropylmethylcellulose 9005-25-8, Starch, biological studies
25322-68-3
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(adhesive containing; buccal delivery of glucagon-like insulinotropic peptides)
- IT 107444-51-9 118549-37-4, Insulinotropin
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(buccal delivery of glucagon-like insulinotropic peptides)


IT 67-68-5, biological studies 68-12-2, biological studies 102-76-1, Triacetin 108-32-7, Propylene carbonate 110-27-0, Isopropyl myristate 111-82-0, Methyl laurate 112-80-1, Oleic acid, biological studies 122-32-7, Glycerol trioleate 127-19-5 143-28-2, Oleyl alcohol 145-42-6, Sodium taurocholate 151-21-3, SDS, biological studies 872-50-4, N-Methylpyrrolidone, biological studies 3445-11-2, N-(2-Hydroxyethyl)-2-pyrrolidinone 5306-85-4, Dimethyl isosorbide 25496-72-4, Glycerol monooleate 25637-84-7, Glycerol dioleate 27194-74-7, Propylene glycol monolaurate 27215-38-9, Glycerol monolaurate 31566-31-1, Glycerol monostearate 59227-89-3
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (permeation enhancer; buccal delivery of glucagon-like insulinotropic peptides)

IT 107-35-7D, Taurine, bile acid conjugates, salts 12441-09-7D, Sorbitan, esters 25312-65-6D, Cholanic acid, salts
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (permeation enhancers; buccal delivery of glucagon-like insulinotropic peptides)

IT 25312-65-6D, Cholanic acid, salts
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (permeation enhancers; buccal delivery of glucagon-like insulinotropic peptides)

RN 25312-65-6 HCAPLUS
 CN Cholan-24-oic acid (8CI, 9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 10 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1997:218959 HCAPLUS
 DN 126:308684
 ED Entered STN: 04 Apr 1997
 TI Use of the intestinal bile acid transporter for the uptake of cholic acid conjugates with HIV-1 protease inhibitory activity
 AU Kagedahle, Matts; Swaan, Peter W.; Redemann, Carl T.; Tang, Mary; Craik, Charles S.; Szoka, Francis C., Jr.; Oie, Svein
 CS Dep. Pharmacy Pharmaceutical Chem., Univ. California, San Francisco, CA, 94143-0446, USA
 SO Pharmaceutical Research (1997), 14(2), 176-180
 CODEN: PHREEB; ISSN: 0724-8741
 PB Plenum
 DT Journal
 LA English
 CC 63-5 (Pharmaceuticals)
 Section cross-reference(s): 1

AB The purpose of this study was to investigate the ability of the human intestinal bile acid transporter to transport cholic acid conjugates with potential HIV-1 protease inhibitory activity. Cholic acid was conjugated at the 24 position of the sterol nucleus with various amino acids and amino acid analogs. The CaCo-2 cell line was used as a model to investigate the interaction of these bile acid conjugates with the human intestinal bile acid transporter. Interaction between the carrier and the conjugates was

quantified by inhibition of taurocholic acid transport and confirmed by transport of radiolabeled conjugates in this cell line. The highest interaction with the transporter, as quantified by inhibition of taurocholic acid transport, occurred when a single neg. charge was present around the 24 to 29 region of the sterol nucleus. A second neg. charge or a pos. charge significantly reduced the interaction. Transport of radiolabeled cholyl-L-Lys- ϵ -tBOC ester and cholyl-D-Asp- β -benzyl ester was inhibited by taurocholic acid. Of all tested compds., only cholyl-D-Asp- β -benzyl ester showed modest HIV-1 protease inhibitory activity with an IC₅₀ of 125 μ M. Cholic acid-amino acid conjugates with appropriate stereochem. are recognized and transported by the human bile acid transporter and show modest HIV-1 protease inhibitory activity. Transport of these conjugates by the bile acid carrier is influenced by charge and hydrophobicity around the 24 position of the sterol nucleus.

ST bile amino acid conjugate intestine transport; HIV1 protease inhibition cholate conjugate AIDS

IT Bile acids

RL: BPR (Biological process); BSU (Biological study, unclassified); PNU (Preparation, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)

(conjugates, with amino acids; use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT Amino acids, biological studies

RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PNU (Preparation, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)

(conjugates, with bile acids; use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT Biological transport

(drug; use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT Drug delivery systems

(oral; use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT Bile acids

RL: BSU (Biological study, unclassified); BIOL (Biological study) (transporter; use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT Biological transport

(uptake; use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT Anti-AIDS agents

Hydrophobicity

Intestine

(use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT 144114-21-6, Retropepsin

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(HIV-1; use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT 7440-23-5, Sodium, biological studies

RL: BSU (Biological study, unclassified); BIOL (Biological study) (bile acid transport dependent on; use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT 2365-14-2P 28071-39-8P, Cholyl-L-lysine

73386-01-3P 89311-00-2P 106335-70-0P

189261-12-9P 189261-13-0P 189261-14-1P

189261-15-2P 189282-94-8P 189282-95-9P

RL: BAC (Biological activity or effector, except adverse); BPR (Biological

process); BSU (Biological study, unclassified); PNU (Preparation, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)

(use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT 81-25-4D, Cholic acid, conjugates with amino acids

RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)

(use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

IT 2365-14-2P 28071-39-8P, Cholyl-L-lysine

73386-01-3P 89311-00-2P 106335-70-0P

189261-12-9P 189261-13-0P 189261-14-1P

189261-15-2P 189282-94-8P 189282-95-9P


RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PNU (Preparation, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)

(use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)

RN 2365-14-2 HCAPLUS

CN L-Cysteine, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 28071-39-8 HCAPLUS

CN L-Lysine, N2-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 73386-01-3 HCAPLUS

CN L-Serine, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 89311-00-2 HCAPLUS

CN D-Alanine, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 106335-70-0 HCAPLUS


CN L-Lysine, N6-[(1,1-dimethylethoxy)carbonyl]-N2-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 189261-12-9 HCAPLUS
 CN D-Alanine, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]-D-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 189261-13-0 HCAPLUS
 CN D-Aspartic acid, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]-, 4-(phenylmethyl) ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 189261-14-1 HCAPLUS
 CN D-Aspartic acid, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]-, 1-(phenylmethyl) ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 189261-15-2 HCAPLUS

CN L-Glutamic acid, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]-, 5-(phenylmethyl) ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 189282-94-8 HCAPLUS

CN D-Aspartic acid, N-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 189282-95-9 HCAPLUS


CN D-Lysine, N6-[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 81-25-4D, Cholic acid, conjugates with amino acids
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
 (use of intestinal bile acid transporter for uptake of cholic acid conjugates with HIV-1 protease inhibitory activity)
 RN 81-25-4 HCAPLUS
 CN Cholan-24-oic acid, 3,7,12-trihydroxy-, (3 α ,5 β ,7 α ,12. α ph a.)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 11 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1995:721131 HCAPLUS
 DN 123:322102
 ED Entered STN: 05 Aug 1995
 TI Acylated derivatives of human insulin with improved solubility and stability for treatment of diabetes
 IN Havelund, Svend; Halstroem, John Broberg; Jonassen, Ib; Andersen, Asser Sloth; Markussen, Jan
 PA Novo Nordisk A/S, Den.
 SO PCT Int. Appl., 99 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM C07K014-62
 ICS A61K038-28
 CC 63-6 (Pharmaceuticals)
 Section cross-reference(s): 2, 3
 FAN.CNT 3

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

Search done by Noble Jarrell

PI WO 9507931 A1 19950323 WO 1994-DK347 19940916 <--
W: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG,
KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU,
SD, SI, SK, TJ, TT, UA, US, UZ, VN
RW: KE, MW, SD, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG
ZA 9407187 A 19950317 ZA 1994-7187 19940916 <--
CA 2171424 AA 19950323 CA 1994-2171424 19940916 <--
CA 2171424 C 20020604
AU 9476520 A1 19950403 AU 1994-76520 19940916 <--
AU 682061 B2 19970918
CN 1133598 A 19961016 CN 1994-193852 19940916 <--
CN 1056618 B 20000920
BR 9407508 A 19970107 BR 1994-7508 19940916 <--
HU 75991 A2 19970528 HU 1996-676 19940916 <--
HU 217684 B 20000328
EP 792290 A1 19970903 EP 1994-926816 19940916 <--
EP 792290 B1 20010829
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE
RO 112873 B1 19980130 RO 1996-583 19940916 <--
JP 3014764 B2 20000228 JP 1995-508923 19940916 <--
JP 09502867 T2 19970325
JP 2000060556 A2 20000229 JP 1999-221632 19940916 <--
PL 178466 B1 20000531 PL 1994-313444 19940916 <--
IL 110977 A1 20000629 IL 1994-110977 19940916 <--
CZ 287945 B6 20010314 CZ 1996-789 19940916 <--
RU 2164520 C2 20010327 RU 1996-108249 19940916 <--
EP 1132404 A2 20010912 EP 2001-112992 19940916 <--
EP 1132404 A3 20020327
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE,
SI, LT
AT 204882 E 20010915 AT 1994-926816 19940916 <--
PT 792290 T 20020130 PT 1994-926816 19940916 <--
ES 2163451 T3 20020201 ES 1994-926816 19940916 <--
SK 282495 B6 20020205 SK 1996-324 19940916 <--
JP 2002308899 A2 20021023 JP 2001-385921 19940916 <--
FI 9601220 A 19960514 FI 1996-1220 19960315 <--
NO 9601070 A 19960515 NO 1996-1070 19960315 <--
AU 9748461 A1 19980219 AU 1997-48461 19971218 <--
AU 745983 B2 20020411 AU 2000-51960 20000811 <--
PRAI DK 1993-1044 A 19930917 <--
US 1994-190829 A 19940202 <--
EP 1994-926816 A3 19940916 <--
JP 1995-508923 A3 19940916 <--
JP 1999-221632 A3 19940916 <--
WO 1994-DK347 W 19940916 <--

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES	
WO 9507931	ICM	C07K014-62	
	ICS	A61K038-28	
WO 9507931	ECLA	C07K014/62	<--
EP 1132404	ECLA	C07K014/62	<--

AB Novel human insulin derivs. with improved solubility and a protracted profile of action are described for use in the treatment of diabetes. These analogs have amino acid substitutions at amino acids A21 and B3 (any amino acid except Lys, Arg, or Cys); PheB1 may be deleted and B30 is substituted by a C10-24 lipophilic amino acid or any naturally occurring amino acid except Lys, Arg, or Cys; if B30 is a lipophilic amino acid, then the ε-NH2 group of LysB29 is acylated with a C≤5 carboxylic acid. They may be used in the treatment of diabetes in several pharmaceutical compns. presented. Chemical preparation of some of these analogs and the manufacture of the amino acid-substituted A and B chains by expression of the cloned cDNAs is demonstrated.

ST human insulin sequence acylation diabetes pharmaceutical
IT Protein sequences

- (acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Solubility
(at physiol. pH; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Acetyl group
Formyl group
(insulin derivs. containing; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Fatty acids, reactions
RL: RCT (Reactant); RACT (Reactant or reagent)
(insulin derivs. containing; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Carboxylic acids, reactions
RL: RCT (Reactant); RACT (Reactant or reagent)
(insulin modification by; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Diabetes mellitus
(insulin pharmaceutical composition for treatment of; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Plasmid and Episome
(pAK-series and pKFN1627 and pEA-series; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Carboxylic acids, reactions
RL: RCT (Reactant); RACT (Reactant or reagent)
(C5, insulins modified with; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Deoxyribonucleic acid sequences
(complementary, acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Carboxylic acids, reactions
RL: RCT (Reactant); RACT (Reactant or reagent)
(di-, C<6, insulin modification by; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Pharmaceutical dosage forms
(injections, insulin; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT Functional groups
(propionyl, insulin derivs. containing; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT 11061-68-ODP, Insulin (human), amino acid-substituted and lipophilic amino acid-containing derivs.
RL: BPN (Biosynthetic preparation); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT 9002-07-7D, Trypsin, immobilized 123175-82-6D, Proteinase, lysine-specific, immobilized
RL: CAT (Catalyst use); PEP (Physical, engineering or chemical process); PROC (Process); USES (Uses)
(acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT 14464-31-4, Palmitic acid N-hydroxysuccinimide ester 69888-86-4
88404-23-3 104943-24-0 165893-02-7 165893-03-8 168986-19-4
168986-20-7 169142-69-2
RL: RCT (Reactant); RACT (Reactant or reagent)
(acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT 168986-17-2P 168986-18-3P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)
- IT 23713-49-7DP, Zn²⁺, complexes with insulin derivs., preparation

RL: SPN (Synthetic preparation); PREP (Preparation)
 (acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 169535-16-4P 169535-18-6P 169535-20-0P 169535-22-2P 169535-28-8P
 169535-30-2P 169535-32-4P 169535-34-6P 169535-36-8P 169535-38-0P
 RL: BPN (Biosynthetic preparation); PRP (Properties); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (amino acid sequence; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 120177-51-7P 169148-61-2P 169148-75-8P
 RL: PRP (Properties); PUR (Purification or recovery); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (amino acid sequence; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 39416-73-4P 169148-55-4DP, zinc complexes 169148-56-5DP, zinc complexes 169148-57-6P 169148-58-7P 169148-59-8P 169148-60-1P
 169148-62-3DP, zinc complexes 169148-63-4P 169148-64-5P 169148-65-6P
 169148-66-7P 169148-67-8P 169148-68-9P 169148-69-0P 169148-70-3P
 169148-71-4P 169148-72-5DP, zinc complexes 169148-72-5P 169148-73-6P
 169148-74-7P
 RL: PRP (Properties); PUR (Purification or recovery); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (amino acid sequence; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 141537-81-7P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (conjugation to insulin; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 168986-14-9P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (for conjugation to insulin; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 7452-59-7, n-Octyl chloroformate
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (in preparation active ester derivs.; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 14565-47-0 22102-66-5 104211-94-1
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (in preparation chemical modified insulin analogs; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 108-30-5, Succinic anhydride, reactions 158627-30-6
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (in preparation myristic acid derivative for conjugation to insulin; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 168986-15-0P 168986-16-1P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (in preparation myristic acid derivative for conjugation to insulin; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 11075-17-5, Carboxypeptidase A
 RL: CAT (Catalyst use); USES (Uses)
 (in preparation of insulin derivs.; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 51-49-0, D-Thyroxine 68528-80-3, Disuccinimidyl suberate
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (in preparation thyroxine derivative for conjugation to insulin; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 110-15-6, Butanedioic acid, reactions 143-07-7, Dodecanoic acid, reactions 638-53-9, Tridecanoic acid 7145-63-3, 2-Aminotetradecanoic

acid 7769-79-1, Hexadecanoic acid, 2-amino- 17702-88-4,
2-Aminodecanoic acid 35237-37-7, 2-Aminododecanoic acid

RL: RCT (Reactant); RACT (Reactant or reagent)

(insulin derivs. containing; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 544-63-8, Tetradecanoic acid, reactions

RL: RCT (Reactant); RACT (Reactant or reagent)

(insulin modification by; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 169535-17-5P 169535-19-7P 169535-21-1P 169535-23-3P 169535-24-4P

169535-25-5P 169535-26-6P 169535-27-7P 169535-29-9P 169535-31-3P

169535-33-5P 169535-35-7P 169535-37-9P 169535-39-1P

RL: BPN (Biosynthetic preparation); PRP (Properties); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(nucleotide sequence; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 24424-99-5, Di-tert-butyl pyrocarbonate

RL: RCT (Reactant); RACT (Reactant or reagent)

(protecting group, in preparation of insulin derivs.; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

IT 168986-19-4


RL: RCT (Reactant); RACT (Reactant or reagent)

(acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

RN 168986-19-4 HCAPLUS

CN Pentanoic acid, 5-[(2,5-dioxo-1-pyrrolidinyl)oxy]-4-[[[(3 α ,5 β)-3-hydroxy-24-oxocholan-24-yl]amino]-5-oxo-, 1,1-dimethylethyl ester, (4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 104211-94-1


RL: RCT (Reactant); RACT (Reactant or reagent)

(in preparation chemical modified insulin analogs; acylated derivs. of human insulin with improved solubility and stability for treatment of diabetes)

RN 104211-94-1 HCAPLUS

CN 2,5-Pyrrolidinedione, 1-[[[(3 α ,5 β)-3-hydroxy-24-oxocholan-24-yl]oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 12 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1995:96130 HCAPLUS
 DN 122:3980
 ED Entered STN: 08 Nov 1994
 TI Bile salts of the toad, *Bufo marinus*: characterization of a new unsaturated higher bile acid, 3 α , 7 α , 12 α , 26-tetrahydroxy-5 β -cholest-23-en-27-oic acid
 AU Yoshii, Michiko; Une, Mizuho; Kihira, Kenji; Kuramoto, Taiju; Akizawa, Toshifumi; Yoshioka, Masanori; Butler, Vincent P., Jr.; Hoshita, Takahiko
 CS Sch. Med., Hiroshima Univ., Hiroshima, 734, Japan
 SO Journal of Lipid Research (1994), 35(9), 1646-51
 CODEN: JLPRAW; ISSN: 0022-2275
 DT Journal
 LA English
 CC 6-5 (General Biochemistry)
 AB The bile salts present in gallbladder bile of the toad, *Bufo marinus*, were found to consist of a mixture of bile alc. sulfates and unconjugated bile acids. The major bile alc. was 5 β -bufol; 5 α - and 5 β -cholestane-3 α , 7 α , 12 α , 26-tetrols occurred as the minor bile alcs. Bile acids of *Bufo marinus* were cholic acid, allocholic acid, 3 α ,7 α ,12 α -trihydroxy-5 α - and 5 β -cholestan-26-oic acids, 3 α ,7 α ,12 α -trihydroxy-5 α - and 5 β -cholest-23-en-26-oic acids, 3 α ,7 α ,12 α ,26-tetrahydroxy-5 β -cholestan-27-oic acid, and a C27 bile acid which has not been previously described. By chromatog. behavior, mass spectral data, and identification of the products of catalytic hydrogenation and ozonolysis, the structure of the new higher bile acid was elucidated as 3 α ,7 α ,12 α ,26-tetrahydroxy-5 β -cholest-23-en-27-oic acid. The bile salt pattern of *Bufo marinus* closely resembles that of *Bufo vulgaris formosus*, except for the absence of 3 α ,7 α ,12 α -trihydroxy-5 β -cholest-22-ene-24-carboxylic acid, the major bile acid of the later toad.
 ST *Bufo* bile salt tetrahydroxycholestenoic acid
 IT Bile
 Bufo marinus
 (bile acids and bile salts of *Bufo marinus*)
 IT Bile acids
 Bile salts
 RL: BOC (Biological occurrence); BSU (Biological study, unclassified);
 BIOL (Biological study); OCCU (Occurrence)
 (bile acids and bile salts of *Bufo marinus*)
 IT 81-25-4, Cholic acid 547-98-8 862-52-2D, 5 α -Cholestane-3 α , 7 α , 12 α , 26-tetrol, sulfate esters 862-53-3D,

5 β -Cholestane-3 α , 7 α , 12 α , 26-tetrol, sulfate esters 2464-18-8 6127-75-9D, sulfate esters 17708-88-2,
3 α , 7 α , 12 α -Trihydroxy-5 α -cholestan-26-oic acid
73834-17-0 84888-63-1 88498-08-2 159330-16-2

RL: BOC (Biological occurrence); BSU (Biological study, unclassified);
BIOL (Biological study); OCCU (Occurrence)
(bile acids and bile salts of *Bufo marinus*)

IT 84888-63-1 88498-08-2


RL: BOC (Biological occurrence); BSU (Biological study, unclassified);
BIOL (Biological study); OCCU (Occurrence)
(bile acids and bile salts of *Bufo marinus*)

RN 84888-63-1 HCAPLUS

CN Cholest-23-en-26-oic acid, 3,7,12-trihydroxy-,
(3 α , 5 β , 7 α , 12 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry unknown.




RN 88498-08-2 HCAPLUS

CN Cholest-23-en-26-oic acid, 3,7,12-trihydroxy-,
(3 α , 5 α , 7 α , 12 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry unknown.



L40 ANSWER 13 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1994:686635 HCAPLUS

DN 121:286635

ED Entered STN: 10 Dec 1994

TI Compositions containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease

IN Sipos, Tibor

PA Digestive Care Inc., USA

SO U.S., 9 pp.

CODEN: USXXAM

DT Patent

LA English
 IC ICM A61K031-56
 INCL 514182000
 CC 63-6 (Pharmaceuticals)
 Section cross-reference(s): 1, 26


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 5352682	A	19941004	US 1993-27693	19930308 <--
PRAI	US 1993-27693		19930308	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 5352682	ICM	A61K031-56
	INCL	514182000
US 5352682	NCL	514/182.000; 424/451.000; 514/788.100; 552/553.000; 552/554.000
	ECLA	A61K031/60

OS MARPAT 121:286635
 GI



AB Disclosed are compns. containing bile acid-aminosalicylate conjugates I (R1 = OH in α or β position; R2 = OH; R3 = H, OH; R4 = H, acetyl) or a pharmaceutically acceptable salt thereof. Also disclosed are a process for preparing the conjugates and methods for treating/preventing gastrointestinal disorders, impaired liver function, etc. using the conjugates.

ST bile acid aminosalicylate conjugate prepn therapeutic; pharmaceutical bile acid aminosalicylate conjugate; deficiency disease bile acid aminosalicylate conjugate; antiinflammatory pharmaceutical bile acid aminosalicylate conjugate

IT Inflammation inhibitors
 Pharmaceutical dosage forms
 (compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)


IT Therapeutics
 (for bile acid deficiency disease; compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)

IT Pharmaceutical dosage forms
 (caplets, compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)

IT Bile acids
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (conjugates, with aminosalicylates; compns. containing acid-aminosalicylate conjugates or salts thereof for

- treating/preventing a bile acid deficiency condition and inflammatory disease)
- IT Bile acids
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (metabolic disorders, deficiency, disease; compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)
- IT Pharmaceutical dosage forms
 (microspheres, compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)
- IT Pharmaceutical dosage forms
 (microtablets, compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)
- IT 28088-64-4DP, Aminosalicyclic acid, bile acid conjugates
 159026-16-1P 159026-17-2P 159026-20-7P
 159026-23-0P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)
- IT 159026-15-0 159026-18-3 159026-19-4
 159026-21-8 159026-22-9 159026-24-1
 159026-25-2 159026-26-3 159026-27-4
 159026-28-5 159026-29-6
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)
- IT 37289-07-9, Cholyglycine hydrolase
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease in relation to conjugate hydrolysis)
- IT 65-49-6, 4-Aminosalicyclic acid 81-25-4, Cholic acid 89-57-6, 5-Aminosalicyclic acid 128-13-2, Ursodeoxycholic acid 474-25-9, Chenodeoxycholic acid
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of and compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)
- IT 159026-16-1P 159026-17-2P 159026-20-7P
 159026-23-0P
 RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (compns. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)
- RN 159026-16-1 HCAPLUS
- CN Benzoic acid, 4-[[(3 α , 5 β , 7 β)-3, 7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-17-2 HCAPLUS

CN Benzoic acid, 5-[[[(3 α ,5 β ,7 β)-3,7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-20-7 HCAPLUS

CN Benzoic acid, 5-[[[(3 α ,5 β ,7 α)-3,7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-23-0 HCAPLUS

CN Benzoic acid, 2-hydroxy-5-[[[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 159026-15-0 159026-18-3 159026-19-4

159026-21-8 159026-22-9 159026-24-1

159026-25-2 159026-26-3 159026-27-4


159026-28-5 159026-29-6

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(comps. containing acid-aminosalicylate conjugates or salts thereof for treating/preventing a bile acid deficiency condition and inflammatory disease)

RN 159026-15-0 HCAPLUS

CN Benzoic acid, 3-[[[(3 α ,5 β ,7 β)-3,7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-18-3 HCAPLUS

CN Benzoic acid, 3-[[[(3 α ,5 β ,7 α)-3,7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-19-4 HCAPLUS

CN Benzoic acid, 4-[[[(3 α ,5 β ,7 α)-3,7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-21-8 HCAPLUS

CN Benzoic acid, 2-hydroxy-3-[[[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-22-9 HCAPLUS

CN Benzoic acid, 2-hydroxy-4-[[[(3 α ,5 β ,7 α ,12 α)-3,7,12-trihydroxy-24-oxocholan-24-yl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-24-1 HCAPLUS

CN Benzoic acid, 2-(acetyloxy)-3-[[[(3 α ,5 β ,12 α)-3,12-dihydroxy-24-oxocholan-24-yl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-25-2 HCAPLUS

CN Benzoic acid, 2-(acetyloxy)-4-[[[(3 α ,5 β ,12 α)-3,12-dihydroxy-24-oxocholan-24-yl]amino]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 159026-26-3 HCAPLUS

CN Benzoic acid, 2-(acetyloxy)-5-[[[(3 α ,5 β ,12 α)-3,12-dihydroxy-24-oxocholan-24-yl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 159026-27-4 HCAPLUS

CN Benzoic acid, 5-[[[(3 α ,5 β ,7 β)-3,7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy-, sodium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 2-A


●x Na

RN 159026-28-5 HCAPLUS

CN Benzoic acid, 5-[[[(3 α ,5 β ,7 β)-3,7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy-, potassium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 2-A

●x K

RN 159026-29-6 HCAPLUS

CN Benzoic acid, 5-[[[(3 α ,5 β ,7 β)-3,7-dihydroxy-24-oxocholan-24-yl]amino]-2-hydroxy-, compd. with 2-amino-2-(hydroxymethyl)-1,3-

Search done by Noble Jarrell


propanediol (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 159026-17-2

CMF C31 H45 N O6


Absolute stereochemistry.



CM 2

CRN 77-86-1

CMF C4 H11 N O3



IT 81-25-4, Cholic acid 128-13-2, Ursodeoxycholic acid

474-25-9, Chenodeoxycholic acid


RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of and compns. containing acid-aminosalicylate conjugates
or salts thereof for treating/preventing a bile acid deficiency
condition and inflammatory disease)

RN 81-25-4 HCAPLUS

CN Cholan-24-oic acid, 3,7,12-trihydroxy-, (3 α ,5 β ,7 α ,12 α .alph
a.)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 128-13-2 HCAPLUS

CN Cholan-24-oic acid, 3,7-dihydroxy-, (3 α ,5 β ,7 β)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 474-25-9 HCAPLUS

CN Cholan-24-oic acid, 3,7-dihydroxy-, (3 α ,5 β ,7 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 14 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1994:173477 HCAPLUS

DN 120:173477

ED Entered STN: 02 Apr 1994

TI The use of nor- and homo- bile acid derivatives as absorption enhancers for medicaments

IN Berlati, Fabio; Ceschel, Giancarlo; Roda, Aldo; Roda, Enrico; Ronchi, Celestino

PA Montereisearch S.r.l., Italy

SO PCT Int. Appl., 14 pp.

Search done by Noble Jarrell

CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM A61K047-28
 CC 63-6 (Pharmaceuticals)
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9400155	A1	19940106	WO 1993-EP1508	19930615 <--
	W: JP, US				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	EP 652773	A1	19950517	EP 1993-912975	19930615 <--
	EP 652773	B1	19980107		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
	JP 07508013	T2	19950907	JP 1993-501998	19930615 <--
	AT 161731	E	19980115	AT 1993-912975	19930615 <--
	ES 2114056	T3	19980516	ES 1993-912975	19930615 <--
	US 5656277	A	19970812	US 1994-360833	19941228 <--
PRAI	IT 1992-MI1601	A	19920630	<--	
	WO 1993-EP1508	W	19930615	<--	

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 9400155	ICM	A61K047-28
US 5656277	NCL	424/400.000; 424/435.000; 424/436.000; 424/451.000; 424/464.000; 424/489.000; 514/169.000; 514/171.000; 514/182.000; 514/553.000; 514/569.000
	ECLA	A61K047/28 <--

AB Nor- and homo- bile acid derivs. and their conjugates with taurine, glycine, and alanine in C23 and C25 are used as absorption enhancers for medicaments administered by the enteral route or by other routes, such as intranasal, buccal and sublingual routes. The derivs. improve the absorption of medicaments through mucosa without being metabolized by the intestinal flora, thus allowing a fast excretion. Moreover, the derivs. have a negligible toxicity. For example, a suppository contained Na diclofenac 0.1, homochenodeoxycholic acid 0.02, and Witepsol H-15 2.5g.

ST bile acid drug absorption enhancer; norbile acid drug absorption enhancer

IT Bile acids
 RL: BIOL (Biological study)
 (as absorption enhancers for drugs)

IT Antihistaminics
 Cardiovascular agents
 Cholinergic antagonists
 Diuretics
 Inflammation inhibitors
 Hormones
 Steroids, biological studies
 RL: BIOL (Biological study)
 (dosage forms of, bile acid derivs. as absorption enhancers in)

IT Peptides, biological studies
 RL: BIOL (Biological study)
 (drugs, dosage forms of, bile acid derivs. as absorption enhancers in)

IT Pharmaceutical dosage forms
 (buccal, bile acid derivs. as absorption enhancers in)

IT Bile acids
 RL: BIOL (Biological study)
 (conjugates, with taurine and glycine and alanine, as absorption enhancers for drugs)


IT Anesthetics
 (local, dosage forms of, bile acid derivs. as absorption enhancers in)

IT Pharmaceutical dosage forms
 (nasal, bile acid derivs. as absorption enhancers in)

IT Bile acids
 RL: BIOL (Biological study)
 (nor-, 3,7-dihydroxy, as absorption enhancers for drugs)


- IT Pharmaceutical dosage forms
(suppositories, bile acid derivs. as absorption enhancers in)
- IT Pharmaceutical dosage forms
(tablets, bile acid derivs. as absorption enhancers in)
- IT 38636-77-0 38636-78-1D, Homochenodeoxycholic acid,
conjugates 53608-86-9, Nordeoxycholic acid
86386-61-0, Norchenodeoxycholic acid 99697-24-2,
Norursodeoxycholic acid 102044-28-0 153311-78-5
153311-79-6 153311-80-9 153481-25-5
RL: BIOL (Biological study)
(as absorption enhancer for drugs)
- IT 56-40-6D, Glycine, conjugates with bile acids 56-41-7D,
Alanine, conjugates with bile acids 107-35-7D, Taurine,
conjugates with bile acids
RL: BIOL (Biological study)
(as absorption enhancers for drugs)
- IT 9034-40-6, LHRH
RL: BIOL (Biological study)
(buccal dosage forms containing, bile acid derivs. as absorption enhancers in)
- IT 9007-12-9, Calcitonin
RL: BIOL (Biological study)
(rectal capsules containing, bile acid derivs. as absorption enhancers in)
- IT 15307-79-6, Sodium diclofenac
RL: BIOL (Biological study)
(suppository containing, bile acid derivs. as absorption enhancers in)
- IT 54-31-9, Furosemide 68-89-3, Dipyrone 443-48-1, Metronidazole
RL: BIOL (Biological study)
(tablets containing, bile acid derivs. as absorption enhancers in)
- IT 38636-77-0 38636-78-1D, Homochenodeoxycholic acid,
conjugates 53608-86-9, Nordeoxycholic acid
86386-61-0, Norchenodeoxycholic acid 99697-24-2,
Norursodeoxycholic acid 102044-28-0 153311-78-5
153311-80-9
RL: BIOL (Biological study)
(as absorption enhancer for drugs)
- RN 38636-77-0 HCAPLUS
- CN Cholan-24-carboxylic acid, 3,12-dihydroxy-, (3 α ,5 β ,12 α)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



- RN 38636-78-1 HCAPLUS
- CN Cholan-24-carboxylic acid, 3,7-dihydroxy-, (3 α ,5 β ,7 α)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 53608-86-9 HCAPLUS

CN 24-Norcholan-23-oic acid, 3,12-dihydroxy-, (3 α ,5 β ,12 α)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 86386-61-0 HCAPLUS

CN 24-Norcholan-23-oic acid, 3,7-dihydroxy-, (3 α ,5 β ,7 α)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 99697-24-2 HCAPLUS

CN 24-Norcholan-23-oic acid, 3,7-dihydroxy-, (3 α ,5 β ,7 β)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 102044-28-0 HCAPLUS

CN Cholan-24-carboxylic acid, 3,7-dihydroxy-, (3 α ,5 β ,7 β)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 153311-78-5 HCAPLUS

CN 24-Norcholan-23-oic acid, 3,12-dihydroxy-, (3 α ,5 β ,12 β)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 153311-80-9 HCAPLUS

CN Cholan-24-carboxylic acid, 3,12-dihydroxy-, (3 α ,5 β ,12 β)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L40 ANSWER 15 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1992:546478 HCAPLUS
 DN 117:146478
 ED Entered STN: 17 Oct 1992
 TI Bile acids and conjugates identified in metabolic disorders by
 fast atom bombardment and tandem mass spectrometry
 AU Libert, Raymond; Hermans, Dominique; Draye, Jean Pierre; Van Hoof,
 Francois; Sokal, Etienne; De Hoffmann, Edmond
 CS Dep. Neuropediatrics, Clin. Univ. St. Luc, Brussels, B-1200, Belg.
 SO Clinical Chemistry (Washington, DC, United States) (1991),
 37(12), 2102-10
 CODEN: CLCHAU; ISSN: 0009-9147
 DT Journal
 LA English
 CC 9-5 (Biochemical Methods)
 Section cross-reference(s): 14, 73, 80
 AB From a study of the collision-activated fragmentation of bile acids, a
 qual. anal. method based on neg.-ion fast-atom-bombardment (FAB) tandem
 mass spectrometry was developed. The times for sample preparation and analyses
 are short. Both free and conjugated bile acids are detected as
 they occur in biol. fluids, acids are detected as they occur in biol.
 fluids, without derivatization. For identifying bile acids and
 conjugates, the method offers better specificity and sensitivity
 than does the fast atom bombardment mass spectrometric technique alone.
 Specific scan modes were developed for the selective detection of taurine
 conjugates, Δ^4 -unsatd. taurine conjugates, Δ^4 -3-keto
 free acids and their glycine conjugates, free
 acids and glycine conjugates bearing a hydroxyl group at the
 C-12 position, sulfates of glycine and taurine conjugates, and a
 C29 dicarboxylic bile acid, specific for generalized peroxisomal
 disorders. Applications of this technique demonstrate its potential
 usefulness, principally in the diagnosis of several peroxisomal disorders.
 ST body fluid bile acid conjugate detection; peroxisome disorder
 diagnosis bile acid; mass spectrometry bile acid diagnosis
 IT Bile
 Blood analysis
 Urine analysis
 (bile acids and their conjugates detection in human, by
 fast-atom-bombardment and tandem mass spectrometry)
 IT Body fluid
 (bile acids and their conjugates detections in, by
 fast-atom-bombardment and tandem mass spectrometry)
 IT Bile acids
 Bile salts
 RL: ANT (Analyte); ANST (Analytical study)
 (detection of, in biol. fluids by fast-atom-bombardment tandem mass
 spectrometry, in metabolic disorder diagnosis)
 IT Mass spectra
 (of bile acids and their conjugates)
 IT Bile acids

RL: ANT (Analyte); ANST (Analytical study)
 (conjugates, detection of, in biol. fluids by
 fast-atom-bombardment tandem mass spectrometry, in metabolic disorder
 diagnosis)

IT Peroxisome
 (disease, diagnosis of, bile acid detection in body fluids by mass
 spectrometry in)

IT Animal metabolism
 (disorder, diagnosis of, bile acid detection in human body fluid by
 mass spectrometry in)

IT Bile acids
 RL: ANT (Analyte); ANST (Analytical study)
 (sulfates, detection of, in biol. fluids by fast-atom-bombardment
 tandem mass spectrometry, in metabolic disorder diagnosis)

IT 516-35-8 640-79-9 13587-11-6 117590-83-7 129944-49-6 143380-61-4
 143380-62-5 143380-63-6 143442-55-1 143476-63-5
 143477-50-3
 RL: ANT (Analyte); ANST (Analytical study)
 (detection of, in biol. fluid by mass spectrometry)

IT 56-40-6, Glycine, analysis 81-24-3, Taurocholic acid 81-25-4
 108-88-3D, Toluene, bile acid conjugates 474-25-9 475-31-0,
 Glycocholic acid 68714-85-2 68756-88-7 117590-89-3 129944-53-2
 143380-64-7 143384-75-2 143442-56-2 143442-57-3 143476-45-3
 143476-62-4
 RL: ANT (Analyte); ANST (Analytical study)
 (detection of, in biol. fluids by mass spectrometry)


IT 60-18-4, L-Tyrosine, analysis
 RL: ANST (Analytical study)
 (metabolic disorders, tyrosine of, type 1, diagnosis of, by bile acid
 mass spectrometry)

IT 143442-55-1
 RL: ANT (Analyte); ANST (Analytical study)
 (detection of, in biol. fluid by mass spectrometry)

RN 143442-55-1 HCAPLUS

CN Cholest-24-en-26-oic acid, 3,7,12-trihydroxy-,
 (3 α ,7 α ,12 α)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry unknown.



L40 ANSWER 16 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1990:421765 HCAPLUS
 DN 113:21765
 ED Entered STN: 21 Jul 1990
 TI Bile acid profiles in peroxisomal 3-oxoacyl-coenzyme A thiolase deficiency
 AU Clayton, Peter T.; Patel, Ella; Lawson, Alexander M.; Carruthers, Robert
 A.; Collins, Janna
 CS Dep. Child Health, Inst. Child Health, London, WC1N 1EH, UK
 SO Journal of Clinical Investigation (1990), 85(4), 1267-73
 CODEN: JCINAO; ISSN: 0021-9738
 DT Journal
 LA English

CC 14-14 (Mammalian Pathological Biochemistry)

AB Fast atom bombardment mass spectrometry and gas chromatog.-mass spectrometry were used to analyze bile acids in the body fluids of an infant (L.C.) whose liver contained no immunoreactive peroxisomal 3-oxoacyl-CoA thiolase. The profiles were compared with those of six patients with undetectable peroxisomes (Zellweger syndrome) and two siblings (N.B. and I.B.) whose defect of peroxisomal β -oxidation could not be localized by morphol. studies of peroxisomes or by immunoblotting of peroxisomal β -oxidation proteins. $3\alpha,7\alpha,12\alpha$ -Trihydroxy- 5β -cholestan-26-oic acid (THCA) was present in bile and plasma of all patients. However, bile from L.C., N.B. and I.B. contained unconjugated varanic acid ($3\alpha,7\alpha,12\alpha,24$ -tetrahydroxy- 5β -cholestan-26-oic acid) as the major C27 bile acid, whereas bile from Zellweger patients contained only small amts. of varanic acid. In the bile from L.C. two isomers of varanic acid were present; in the bile from N.B. and I.B. a single isomer predominated. L.C., N.B., and I.B. all produced bile containing small amts. of (24E)- $3\alpha,7\alpha,12\alpha$ -trihydroxy- 5β -cholest-24-en-26-oic acid ([24E]- Δ 24-THCA), its [24Z]-isomer, $3\alpha,7\alpha,12\alpha$ -trihydroxy- 5β -cholest-23-en-26-oic acid and $3\alpha,7\alpha,12\alpha$ -trihydroxy-27-nor- 5β -cholestan-24-one. The results provide evidence for peroxisomal pathways for cholic acid synthesis in man via THCA, Δ 24-THCA, and varanic acid and show that bile acid analyses can be used to diagnose peroxisomal thiolase deficiency.

ST bile acid profile oxoacylCoA thiolase deficiency

IT Blood plasma

Urine
(bile acid profiles of, in peroxisomal oxoacyl-CoA thiolase deficiency of humans, diagnosis in relation to)

IT Body fluid
(duodenal, bile acid profiles of, in diagnosis of peroxisomal oxoacyl-CoA thiolase deficiency of humans)

IT Bile acids
RL: BIOL (Biological study)
(of body fluids, in peroxisomal oxoacyl-CoA thiolase deficiency of humans, diagnosis in relation to)

IT Peroxisome
(oxoacyl-CoA thiolase deficiency of, bile acid profiles of body fluids in diagnosis of, of humans)

IT 9029-97-4, 3-Oxoacyl-CoA thiolase
RL: BIOL (Biological study)
(deficiency of, bile acid profiles of duodenal juice in diagnosis of, of humans)


IT 81-25-4, Cholic acid 128-13-2, Ursodeoxycholic acid 474-25-9, Chenodeoxycholic acid 547-98-8 1061-64-9 61628-32-8 72883-89-7 73834-17-0 84888-63-1 85552-38-1 85552-39-2 85552-42-7
RL: BIOL (Biological study)
(of duodenal juice, in peroxisomal oxoacyl-CoA thiolase deficiency of humans, diagnosis in relation to)

IT 84888-63-1 85552-38-1 85552-39-2
RL: BIOL (Biological study)
(of duodenal juice, in peroxisomal oxoacyl-CoA thiolase deficiency of humans, diagnosis in relation to)

RN 84888-63-1 HCAPLUS


CN Cholest-23-en-26-oic acid, 3,7,12-trihydroxy-,
($3\alpha,5\beta,7\alpha,12\alpha$)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry unknown.




RN 85552-38-1 HCAPLUS
 CN Cholest-24-en-26-oic acid, 3,7,12-trihydroxy-,
 (3 α ,5 β ,7 α ,12 α ,24E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.




RN 85552-39-2 HCAPLUS
 CN Cholest-24-en-26-oic acid, 3,7,12-trihydroxy-,
 (3 α ,5 β ,7 α ,12 α ,24Z)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.



L40 ANSWER 17 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1988:471021 HCAPLUS
 DN 109:71021
 ED Entered STN: 02 Sep 1988
 TI Occurrence of 3 β -hydroxy-5-cholestenoic acid, 3 β ,7 α -
 dihydroxy-5-cholestenoic acid, and 7 α -hydroxy-3-oxo-4-cholestenoic
 acid as normal constituents in human blood
 AU Axelson, Magnus; Moerk, Birgitta; Sjoevall, Jan

CS Dep. Clin. Chem., Karolinska Hosp., Stockholm, 104 01, Swed.
 SO Journal of Lipid Research (1988), 29(5), 629-41
 CODEN: JLPRAW; ISSN: 0022-2275
 DT Journal
 LA English
 CC 13-5 (Mammalian Biochemistry)
 GI




I, $R=R^2=H$, $R^1=OH$
 II, $R=H$, $R^1=R^2=OH$
 III, $RR^1=O$, $R^2=OH$


AB Three unconjugated C27 bile acids were found in plasma from healthy humans. They were isolated by liquid-solid extraction and anion-exchange chromatog. and were identified by gas-liquid chromatog.-mass spectrometry, microchem. reactions, and UV spectroscopy as 3β -hydroxy-5-cholestenoic, $3\beta,7\alpha$ -dihydroxy-5-cholestenoic, and 7α -hydroxy-3-oxo-4-cholestenoic acids (I, II, and III, resp.). Their levels often exceeded those of the unconjugated C24 bile acids and the variations between individuals were smaller than for the C24 acids. The concns. in plasma from healthy subjects were 67.2 ng/mL for I, 38.9 ng/mL for II, and 81.7 ng/mL for III. The levels of the individual acids were pos. correlated with each other and not with the levels of the C24 acids. The cholestenic acids were below the detection limit (20-50 ng/mL) in bile, and C27 bile acids present in bile were not detected in plasma.

ST cholestenic acid deriv blood plasma; bile acid C27 blood plasma
 IT Feeding
 (bile acids of blood plasma of human response to)
 IT Bile
 Blood plasma
 (bile acids of, of human)
 IT Bile acids
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (of bile, of human, bile acids of blood plasma in relation to)
 IT Bile acids
 RL: BIOL (Biological study)
 (of blood plasma of human)
 IT 81-25-4, Cholic acid 474-25-9, Chenodeoxycholic acid 547-98-8
 5226-26-6 60696-62-0, Norcholeic acid 72883-89-7 73804-37-2
 73834-17-0, $3\alpha,7\alpha,12\alpha,26$ -Tetrahydroxy-5 β -cholestan-27-oic acid 73837-07-7
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (of bile, of human)
 IT 6561-58-6 115538-84-6 115538-85-7
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (of blood plasma, of human)
 IT 115567-29-8
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (oxidation of)
 IT 115538-86-8P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)

Absolute stereochemistry.
Double bond geometry unknown.



L40 ANSWER 18 OF 19 HCAPLUS COPYRIGHT 2005 ACS on STN
AN 1988:187048 HCAPLUS
DN 108:187048
ED Entered STN: 28 May 1988
TI Bile acids with a cyclopropyl-containing side chain. 3. Separation,
identification, and properties of all four stereoisomers of
3 α ,7 β -dihydroxy-22,23-methylene-5 β -cholan-24-oic acid
AU Pellicciari, Roberto; Natalini, Benedetto; Cecchetti, Sergio; Porter,
Barry; Roda, Aldo; Grigolo, Brunella; Balducci, Renzo
CS Ist. Chim. Farm. Tec. Farm., Univ. Studi, Perugia, 06100, Italy
SO Journal of Medicinal Chemistry (1988), 31(4), 730-6
CODEN: JMCMAR; ISSN: 0022-2623.
DT Journal
LA English
CC 32-6 (Steroids)
OS CASREACT 108:187048
GI




I

Search done by Noble Jarrell

concentration in water, and exhibit a different interaction with intestinal bacterial enzymes. CUDCA A-C are not conjugated with glycine or taurine in the liver, while CUDCA D is secreted into bile predominantly as taurine and glycine conjugate.

- ST methylencholanoic acid stereoisomer configuration; cholanic acid methylene stereoisomer configuration; bile acid biol activity diastereoisomer
- IT Molecular structure-biological activity relationship
(of 3 α , 7 β -dihydroxy-22,23-methylene-5 β -cholan-24-oic acid diastereomers)
- IT 105360-63-2
RL: RCT (Reactant); RACT (Reactant or reagent)
(cyclopropanation of, with Et diazoacetate)
- IT 128-13-2
RL: RCT (Reactant); RACT (Reactant or reagent)
(esterification of)
- IT 113181-05-8P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and acetylation of)
- IT 113218-62-5P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and addition reaction with Et diazoacetate)
- IT 113181-08-1P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and attempted cyclopropanation of)
- IT 113299-41-5P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and conversion into acetylene derivative)
- IT 113181-06-9P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and cyclopropanation with diazomethane)
- IT 113181-04-7P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and elimination reaction of)
- IT 113181-07-0P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and hydrogenation of)
- IT 69519-36-4P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and phenylselenylation of)
- IT 89414-90-4P 89495-32-9P 89495-33-0P 89495-34-1P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and saponification of)
- IT 91378-92-6P 91423-31-3P 91423-32-4P 91423-33-5P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation, purification, configuration, and biol. activity of)
- IT 113181-05-8P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and acetylation of)
- RN 113181-05-8 HCAPLUS
- CN Chol-22-en-24-oic acid, 3,7-dihydroxy-, ethyl ester,
(3 α , 5 β , 7 β , 22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



```

L40  ANSWER 19 OF 19 HCAPLUS  COPYRIGHT 2005 ACS on STN
AN   1987:154139  HCAPLUS
DN   106:154139
ED   Entered STN: 15 May 1987
TI   Identification of unconjugated bile acids in human bile
AU   Matoba, Naoyuki; Une, Mizuho; Hoshita, Takahiko
CS   Fac. Med., Kyushu Univ., Maidashi, 3-1-1, Japan
SO   Journal of Lipid Research (1986), 27(11), 1154-62
     CODEN: JLPRAW; ISSN: 0022-2275
DT   Journal
LA   English
CC   14-7 (Mammalian Pathological Biochemistry)
     Section cross-reference(s): 13
AB   Unconjugated bile acids in the bile of healthy and diseased
     (cerebrotendinous xanthomatosis) humans were determined qual. and quant. by
     gas-liquid chromatog. and gas-liquid chromatog.-mass spectrometry, after their
     isolation by ion-exchange chromatog. In a healthy person and 3 patients
     with cholelithiasis, unconjugated bile acids comprised 0.1-0.4%
     of total biliary bile acids. The bile acid composition of the
     unconjugated fraction was quite different from that of the
     glycine- or taurine-conjugate fraction, in that it contained a
     relatively large proportion of unusual bile acids including C23 and C27
     bile acids. In 2 patients with cerebrotendinous xanthomatosis, C22 and
     C23 bile acids were the major constituents of the biliary
     unconjugated bile acids and comprised about 0.8% of total bile
     acids; no detectable amts. of C27 bile acids were found in their bile.
     The anal. of biliary unconjugated bile acids may be useful for
     the diagnosis of metabolic diseases concerning bile acids, particularly
     those diseases which involve the accumulation or disappearance of unusual
     bile acids.
ST   bile acid bile cholelithiasis cerebrotendinous xanthomatosis
IT   Calculi, biliary
     (unconjugated bile acids of bile in, in humans)
IT   Bile
     (unconjugated bile acids of, of humans)
IT   Bile acids
     RL: BIOL (Biological study)
     (unconjugated, of bile in cerebrotendinous xanthomatosis and
     cholelithiasis and health in humans)
IT   Xanthomatosis
     (cerebrotendinous, unconjugated bile acids of bile in, in
     humans)
IT   56-40-6, biological studies 107-35-7, Taurine
     RL: BIOL (Biological study)
     (bile acids conjugated with, of bile in cerebrotendinous
     xanthomatosis and cholelithiasis and health in humans)
IT   81-25-4, Cholic acid 83-44-3, Deoxycholic acid 128-13-2,
     Ursodeoxycholic acid 474-25-9, Chenodeoxycholic acid 547-98-8

```

911-40-0, 7-Ketodeoxycholic acid 2464-18-8, Allocholic acid 2955-27-3,
 7-Epicholic acid 38917-20-3 60696-62-0, Norcholic acid 61844-74-4
 73804-37-2 73834-17-0 73837-07-7 85552-39-2 86386-61-0
 98349-18-9 99697-24-2 107480-95-5

RL: BIOL (Biological study)

(unconjugated, of bile in cerebrotendinous xanthomatosis and
 cholelithiasis and health in humans)

IT 85552-39-2 107480-95-5

RL: BIOL (Biological study)


(unconjugated, of bile in cerebrotendinous xanthomatosis and
 cholelithiasis and health in humans)

RN 85552-39-2 HCAPLUS

CN Cholest-24-en-26-oic acid, 3,7,12-trihydroxy-,
 (3 α ,5 β ,7 α ,12 α ,24Z)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




RN 107480-95-5 HCAPLUS

CN Chol-22-en-24-oic acid, 3,7,12-trihydroxy-, (3 α ,5 β ,7 α ,12.
 alpha.,22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.



=> b embase

FILE 'EMBASE' ENTERED AT 09:39:11 ON 24 JUN 2005
 COPYRIGHT (C) 2005 Elsevier Inc. All rights reserved.

FILE COVERS 1974 TO 23 Jun 2005 (20050623/ED)

EMBASE has been reloaded. Enter HELP RLOAD for details.

This file contains CAS Registry Numbers for easy and accurate
 substance identification.

Search done by Noble Jarrell

=> d all 148 tot

L48 ANSWER 1 OF 4 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN
AN 2004197875 EMBASE
TI Absorption of the cholic acid-conjugated peptide hormone cholylsecretin
from the rat ileum in vivo.
AU McHarg S.; Morton J.S.; McGinn B.J.; Yasin M.; Morrison J.D.
CS J.D. Morrison, West Medical Building, University of Glasgow, Glasgow G12
8QQ, United Kingdom
SO Acta Physiologica Scandinavica, (2004) Vol. 181, No. 1, pp. 23-34.
Refs: 29
ISSN: 0001-6772 CODEN: APSCAX
CY United Kingdom
DT Journal; Article
FS 003 Endocrinology
030 Pharmacology
037 Drug Literature Index
048 Gastroenterology
LA English
SL English
ED Entered STN: 20040610
Last Updated on STN: 20040610
AB Aims: Previously, we demonstrated that gastrin peptides as long as 34
amino acids were absorbed from the ileum of rat after conjugation to the
C24 position of cholic acid and that these peptides retained full
biological activity. As absorption was specific to the ileum, it was
inferred that the conjugated hormone was taken up by the bile salt
transporters. We have now extended these experiments to a member of a
different family of hormones, viz. secretin, a 27-amino acid hormone that
stimulates serous secretions from the exocrine pancreas. Methods: After
conjugation to cholic acid, the degree of cholylsecretin absorption from
the ileum of anaesthetized rats was assessed from the increase in
pancreatic secretions. Results: A complication to the study was that
intra-ileal infusion of native secretin caused a transient increase in the
levels of pancreatic secretions. This was in contrast to the effects of
intra-ileal infusion of cholylsecretin which did not cause this transient
increase but, instead, gave rise to a delayed increase in pancreatic
secretions which was sustained over several hours during which
cholylsecretin was detected in plasma in high concentration by mass
spectrometry. The pancreatic response to cholylsecretin was abolished by
co-infusion of 50 mM taurocholate, employed to compete with the bile salt
transporters, although a transient increase in pancreatic secretions
similar to that caused by secretin was now generated. This was shown to
arise from an action of taurocholate per se causing the release of
endogenous secretin which is present in rat ileum. Conclusions: We,
therefore, concluded that cholylsecretin had been absorbed from the rat
ileum by uptake by bile salt transporters.
CT Medical Descriptors:
*hormone release
*hormonal regulation
*small intestine absorption
*ileum
drug effect
drug efficacy
drug mechanism
pancreas secretion
hormone blood level
secretin blood level
intestine absorption
pharmacological blocking
pancreas
nonhuman
male
rat

animal experiment
 controlled study
 animal tissue
 article
 priority journal
 Drug Descriptors:
 *cholic acid: PD, pharmacology
 *secretin: EC, endogenous compound
 *secretin: PD, pharmacology
 recombinant hormone: PD, pharmacology
 carrier protein: EC, endogenous compound
 sodium chloride: EC, endogenous compound
 taurocholic acid: PD, pharmacology
 RN (cholic acid) 32500-01-9, 361-09-1, 81-25-4;
 (secretin) 1393-25-5, 17034-35-4, 73559-81-6; (carrier protein)
 80700-39-6; (sodium chloride) 7647-14-5; (taurocholic acid) 145-42-6,
 59005-70-8, 81-24-3
 CO Sigma Aldrich

 L48 ANSWER 2 OF 4 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 2002415967 EMBASE
 TI Absorption of biologically active peptide hormones from the small
 intestine of rat.
 AU Wheeler S.; McGinn B.J.; Lucas M.L.; Morrison
 J.D.
 CS J.D. Morrison, West Medical Building, University of Glasgow, Glasgow G12
 8QQ, United Kingdom
 SO Acta Physiologica Scandinavica, (2002) Vol. 176, No. 3, pp. 203-213.
 Refs: 34
 ISSN: 0001-6772 CODEN: APSCAX
 CY United Kingdom
 DT Journal; Article
 FS 002 Physiology
 030 Pharmacology
 037 Drug Literature Index
 048 Gastroenterology
 LA English
 SL English
 ED Entered STN: 20021202
 Last Updated on STN: 20021202
 AB Absorption of the 4, 10 and 34 amino acid forms of gastrin from the small
 intestine has been investigated in anaesthetized rats. The method of
 assessment of successful absorption of the hormone into the systemic
 circulation was when the amount of acid secreted by the stomach over
 consecutive 15-min periods was increased. When the natural hormones were
 infused into the ileum in a relatively high dose, there was no increase in
 gastric acid secretion, indicating that they had not been absorbed. Each
 of the forms of gastrin was conjugated at the free amino terminus to the
 carboxyl group of cholic acid. Subsequent infusion of the conjugated form
 of gastrin into the ileum, this time in relatively low doses, resulted in
 substantial and prolonged increases in gastric acid secretion, indicating
 that these hormones had been successfully absorbed. In addition,
 conjugation of the 10 and 34 amino acid forms of gastrin with cholic acid
 was shown to increase markedly the potency in evoking an increase in
 gastric acid secretion in response to intravenous injection of the
 hormone. Absorption of the gastrin conjugates was specific to the ileum
 thus indicating that they had been absorbed through the bile salt
 transporters.
 CT Medical Descriptors:
 *small intestine absorption
 systemic circulation
 acid secretion
 ileum
 stomach acid secretion
 amino terminal sequence

conjugation
 carboxy terminal sequence
 drug effect
 drug megadose
 nonhuman

male
 rat
 animal experiment
 controlled study
 animal tissue

article
 priority journal
 Drug Descriptors:

*peptide hormone: DO, drug dose
 *peptide hormone: PD, pharmacology
 *peptide hormone: IV, intravenous drug administration
 *gastrin: DO, drug dose
 *gastrin: PD, pharmacology
 *gastrin: IV, intravenous drug administration
 cholic acid

RN (gastrin) 9002-76-0; (cholic acid) 32500-01-9, 361-09-1
 , 81-25-4

L48 ANSWER 3 OF 4 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN

AN 91218502 EMBASE

DN 1991218502

TI The effect of sodium deoxycholate and other surfactants on the mucosal
 surface pH in proximal jejunum or rat.

AU McKie A.T.; Stewart W.; Lucas M.L.

CS Institute of Physiology, Glasgow University, Glasgow G12 8QQ, United
 Kingdom

SO Naunyn-Schmiedeberg's Archives of Pharmacology, (1991) Vol. 343, No. 6,
 pp. 659-664.

ISSN: 0028-1298 CODEN: NSAPCC

CY Germany

DT Journal; Article

FS 002 Physiology
 004 Microbiology
 029 Clinical Biochemistry
 048 Gastroenterology
 052 Toxicology
 030 Pharmacology
 037 Drug Literature Index

LA English

SL English

ED Entered STN: 911216

Last Updated on STN: 911216

AB The mucosal surface pH (acid microclimate) and nucleotide levels of rat
 proximal jejunum were measured in vivo under various conditions which
 included exposure to pharmacological agents and to surfactants. Mucosal
 surface pH was unaffected by sodium nitroprusside, A23187 and amiloride,
 as was mucosal cGMP content, although amiloride and A23187 reduced cAMP
 content. In contrast, surfactants elevated the pH of the mucosal surface
 significantly ($P < 0.001$): control value 6.23 ± 0.02 ($n = 12$); Lubrol
 PX 0.8% (v/v) 6.98 ± 0.02 ($n = 5$); sodium deoxycholate 2 mmol/l 6.67
 ± 0.04 ($n = 5$); Triton X-100 0.5% (v/v) 7.41 ± 0.03 ($n = 5$). No
 significant changes in cGMP levels were noted after surfactant treatment,
 although DOC and Triton X-100 reduced cAMP levels. The ability of higher
 concentrations of surfactant to elevate the mucosal surface pH beyond
 neutrality to values similar to plasma pH contrasts with the action of
 Escherichia coli heat-stable (STa) enterotoxin which at high
 concentrations could not elevate the mucosal surface pH beyond neutrality.
 Consistent with the known effects on tight junction permeability,
 surfactants may act by allowing plasma-like subepithelial fluid to
 neutralise the microclimate.

CT Medical Descriptors:
 *cell surface
 *jejunum mucosa
 *ph
 animal experiment
 animal tissue
 article
 controlled study
 male
 microscopy
 nonhuman
 priority journal
 radioimmunoassay
 rat
 regional perfusion
 Drug Descriptors:
 *amiloride: PD, pharmacology
 *calcimycin: PD, pharmacology
 *cyclic gmp: EC, endogenous compound
 *deoxycholate sodium: TO, drug toxicity
 *nitroprusside sodium: PD, pharmacology
 *surfactant: TO, drug toxicity
 cyclic amp: EC, endogenous compound
 docusate sodium: TO, drug toxicity
 escherichia coli enterotoxin: TO, drug toxicity
 lubrol: TO, drug toxicity
 triton x 100: TO, drug toxicity
 RN (amiloride) 2016-88-8, 2609-46-3; (calcimycin) 52665-69-7; (cyclic gmp) 7665-99-8; (deoxycholate sodium) 302-95-4; (nitroprusside sodium) 14402-89-2, 15078-28-1; (cyclic amp) 60-92-4; (docusate sodium) 577-11-7; (lubrol) 11138-41-3, 52434-01-2
 CN Lubrol px; Triton x 100; A 23187
 L48 ANSWER 4 OF 4 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 83183463 EMBASE
 DN 1983183463
 TI The effect of deoxycholate on intestinal surface pH and 5-methyltetrahydropteroylglutamate absorption in the rat proximal jejunum in vitro.
 AU Blair J.A.; Hilburn M.E.; Lucas M.L.; Said H.M.
 CS Dep. Chem., Univ. Aston Birmingham, Birmingham B4 7ET, United Kingdom
 SQ Biochemical Society Transactions, (1983) Vol. 11, No. 2, pp. 165-167.
 CODEN: BCSTB5
 CY United Kingdom
 DT Journal
 FS 037 Drug Literature Index
 029 Clinical Biochemistry
 002 Physiology
 048 Gastroenterology
 LA English
 ED Entered STN: 911209
 Last Updated on STN: 911209
 CT Medical Descriptors:
 *5 methyltetrahydrofolic acid c 14
 *drug absorption
 *intestine absorption
 *intestine mucosa
 *ph
 jejunum
 nonhuman
 rat
 small intestine
 animal cell
 digestive system
 Drug Descriptors:

*5 methyltetrahydrofolic acid
 *deoxycholic acid
 radioisotope
 RN (5 methyltetrahydrofolic acid) 134-35-0; (deoxycholic acid)
 83-44-3

=> d all 151 tot

L51 ANSWER 1 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 1999324376 EMBASE
 TI Study of the pharmacological effect of the bile salt, sodium scymnol
 sulfate, from Rhizoprionodon acutus. IV. Effects of naturally occurring
 bile alcohols, bile acids and their conjugates on lesion
 development and vascular endothelial cell injury in a rat peripheral
 arterial occlusion model.
 AU Ishida H.; Nakayasu H.; Tsuji K.
 CS H. Ishida, School of Pharmaceutical Science, University of Shizuoka, 52-1
 Yada, Shizuoka 422-8526, Japan
 SO Biological and Pharmaceutical Bulletin, (1999) Vol. 22, No. 8, pp.
 828-835.
 Refs: 48
 ISSN: 0918-6158 CODEN: BPBLEO
 CY Japan
 DT Journal; Article
 FS 018 Cardiovascular Diseases and Cardiovascular Surgery
 030 Pharmacology
 037 Drug Literature Index
 LA English
 SL English
 ED Entered STN: 19990930
 Last Updated on STN: 19990930
 AB A series of naturally occurring bile alcohols, bile acids and their
 conjugates has been investigated as part of our studies to develop
 unique anticoagulants with a potent prophylactic effect against vascular
 endothelial cell injury induced by lactic acidosis in vivo and in vitro.
 In an in vivo rat peripheral arterial occlusion model induced by lactic
 acid injection, oral administration of a single dose of 3 mg/kg scymnol
 significantly inhibited edematous swelling and development of lower limb
 lesions, including gangrene, and reduced changes in clotting system
 functions and serum lactate dehydrogenase activity. It had no effect on
 clotting system functions in sham-operated rats. The structure-activity
 relationship suggests that the [24R-(+)-5 β -cholestane-
 3 α ,7 α ,24,26-pentol] or [3 α ,7 α -dihydroxy-5 β -
 cholanolic acid] structure is important for a potent prophylactic effect
 following oral administration. Intravenous administration of a single
 dose of 0.3 mg/kg sodium (25S)-scymnol sulfate or scymnol prevented lesion
 progression as effectively as oral administration of scymnol. Sodium
 (25S)-scymnol sulfate and ursodeoxycholic acid showed clear protective
 effects against cultured vascular endothelial cell damage due to lactic
 acidosis which were dose-dependent. The above results suggest that bile
 steroids such as scymnol, sodium (25S)-scymnol sulfate, ursodeoxycholic
 acid, and chenodeoxycholic acid may play a role in protecting endothelial
 cells against injury caused by lactic acidosis. These compounds are
 candidates for novel anti-ischemic drugs that act by specifically
 protecting vascular endothelial cells.
 CT Medical Descriptors:
 *peripheral occlusive artery disease: DT, drug therapy
 *peripheral occlusive artery disease: PC, prevention
 gangrene: PC, prevention
 structure activity relation
 lactate dehydrogenase blood level
 prophylaxis
 vascular endothelium

cell protection
 nonhuman
 male
 rat
 animal experiment
 animal model
 controlled study
 animal tissue
 animal cell
 oral drug administration
 intravenous drug administration
 intraperitoneal drug administration
 article
 Drug Descriptors:
 *bile salt: DT, drug therapy
 *bile acid: DT, drug therapy
 *bile acid conjugate: DT, drug therapy
 *scymnol: DT, drug therapy
 lactic acid
 lactate dehydrogenase: EC, endogenous compound
 ursodeoxycholic acid: DT, drug therapy
 argatroban: DT, drug therapy
 chenodeoxycholic acid: DT, drug therapy
 cholic acid: DT, drug therapy
 tauroursodeoxycholic acid: DT, drug therapy
 taurochenodeoxycholic acid: DT, drug therapy
 taurocholic acid: DT, drug therapy
 RN (lactic acid) 113-21-3, 50-21-5; (lactate dehydrogenase) 9001-60-9;
 (ursodeoxycholic acid) 128-13-2, 2898-95-5;
 (argatroban) 74863-84-6; (chenodeoxycholic acid) 474-25-9;
 (cholic acid) 32500-01-9, 361-09-1, 81-25-4;
 (tauroursodeoxycholic acid) 14605-22-2; (taurochenodeoxycholic acid)
 516-35-8; (taurocholic acid) 145-42-6, 59005-70-8, 81-24-3
 CO Wako; Tokyo Tanabe; Sigma; Daiichi Pharmaceutical (Japan)
 L51 ANSWER 2 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 1999256785 EMBASE
 TI Simultaneous determination of ursodeoxycholic acid and its glycine-
 conjugate in serum as phenacyl esters using multidimensional
 liquid chromatography.
 AU Choi S.J.; Jeong C.K.; Lee H.M.; Kim K.; Do K.S.; Lee H.S.
 CS S.J. Choi, College of Pharmacy, Wonkwang University, Iksan 570-749, Korea,
 Republic of
 SO Chromatographia, (1999) Vol. 50, No. 1-2, pp. 96-100.
 Refs: 25
 ISSN: 0009-5893 CODEN: CHRGB7
 CY Germany
 DT Journal; Article
 FS 030 Pharmacology
 037 Drug Literature Index
 LA English
 SL English
 ED Entered STN: 19990812
 Last Updated on STN: 19990812
 AB A narrowbore high-performance liquid chromatographic (HPLC) method using
 column switching is described for the simultaneous determination of
 ursodeoxycholic acid (UDCA) and glyco-UDCA (GUDCA) from serum samples as
 their phenacyl esters. Serum samples were subjected to a preliminary
 clean-up using octadecylsilane reversed-phase extraction and derivatized
 with phenacylbromide. The purification, fractionation and concentration
 of UDCA and GUDCA from the esterified serum sample were performed on-line
 by appropriate switching of columns. Limit of detection (LOD) of UDCA and
 GUDCA were 5 ng and the absolute mean recoveries averaged 84.4 ± 8.2%
 and 85.2 ± 8.4%, respectively. This method was successfully applied
 to the pharmacokinetic study of UDCA in rats and human.

CT Medical Descriptors:
 *high performance liquid chromatography
 extraction
 purification
 fractionation
 drug blood level
 validation process
 human
 nonhuman
 rat
 oral drug administration
 intravenous drug administration
 article
 priority journal
 Drug Descriptors:
 *ursodeoxycholic acid: AD, drug administration
 *ursodeoxycholic acid: CR, drug concentration
 *ursodeoxycholic acid: DO, drug dose
 *ursodeoxycholic acid: PK, pharmacokinetics
 *glycine
 *ester derivative
 *glycoursodeoxycholic acid
 silane derivative
 bromine derivative
 RN (ursodeoxycholic acid) 128-13-2, 2898-95-5; (glycine)
 56-40-6, 6000-43-7, 6000-44-8; (glycoursodeoxycholic acid)
 64480-66-6
 CO Sigma (United States)

L51 ANSWER 3 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 1999046272 EMBASE
 TI Inhibition of protein denaturation by fatty acids, bile salts and other
 natural substances: A new hypothesis for the mechanism of action of fish
 oil in rheumatic diseases.
 AU Saso L.; Valentini G.; Casini M.L.; Mattei E.; Braghiroli L.; Mazzanti G.;
 Panzironi C.; Grippa E.; Silvestrini B.
 CS B. Silvestrini, Inst. Pharmacology and Pharmacognosy, University 'La
 Sapienza', P.le Aldo Moro 5, 00185 Rome, Italy
 SO Japanese Journal of Pharmacology, (1999) Vol. 79, No. 1, pp. 89-99.
 Refs: 41
 ISSN: 0021-5198 CODEN: JJPAAZ
 CY Japan
 DT Journal; Article
 FS 017 Public Health, Social Medicine and Epidemiology
 026 Immunology, Serology and Transplantation
 029 Clinical Biochemistry
 030 Pharmacology
 031 Arthritis and Rheumatism
 037 Drug Literature Index
 LA English
 SL English
 ED Entered STN: 19990218
 Last Updated on STN: 19990218
 AB Natural hydrophobic substances like bile salts (cholate, deoxycholate,
 chenodeoxycholate, lithocholate and their conjugates with
 glycine and taurine), fatty acids (caprylic, capric, lauric, myristic,
 palmitic, stearic, oleic, linoleic, arachidonic, eicosapentaenoic and
 docosahexaenoic acid) were much more active (EC50 .simeq. 10-4-10-5 M)
 than selected amino acids (EC50 > 10-2 M) and inorganic salts (EC50
 .simeq. 10-1 M) in inhibiting heat-induced denaturation of human serum
 albumin in vitro. Fish oil, rich in n-3-polyunsaturated acids such as
 eicosapentaenoic acid and docosahexaenoic acid, administered p.o. (1
 ml/kg) in the rat, protected ex vivo (after 2 hr) serum against
 heat-induced denaturation more than bendazac, a known antidenaturant drug.
 Thus, we speculated that the antidenaturant activity of fish oil may be

partly (in addition to the known effect on endogenous eicosanoid composition) responsible for its beneficial effects in rheumatoid arthritis and other rheumatic conditions. In this connection, it is of note that the in vitro antidenaturant activity of fish oil fatty acids was higher than that of known antidenaturant drugs such as bendazac and bindarit and nonsteroidal anti-inflammatory drugs like phenylbutazone and indomethacin which could exert beneficial effects in chronic inflammatory conditions by stabilizing endogenous proteins.

CT

Medical Descriptors:

- *protein denaturation
- *drug mechanism
- *rheumatic disease: DT, drug therapy
- protein stability
- cattle
- human
- nonhuman
- rat
- normal human
- animal experiment
- controlled study
- human tissue
- animal tissue

- oral drug administration

article

Drug Descriptors:

- *fatty acid: DV, drug development
- *fatty acid: PD, pharmacology
- *fish oil: DV, drug development
- *fish oil: DT, drug therapy
- *fish oil: PD, pharmacology
- *bile salt: DV, drug development
- *bile salt: PD, pharmacology
- cholic acid: CM, drug comparison
- cholic acid: DV, drug development
- cholic acid: PD, pharmacology
- deoxycholic acid: CM, drug comparison
- deoxycholic acid: DV, drug development
- deoxycholic acid: PD, pharmacology
- chenodeoxycholic acid: CM, drug comparison
- chenodeoxycholic acid: DV, drug development
- chenodeoxycholic acid: PD, pharmacology
- lithocholic acid: CM, drug comparison
- lithocholic acid: DV, drug development
- lithocholic acid: PD, pharmacology
- bile acid conjugate: CM, drug comparison
 - bile acid conjugate: DV, drug development
 - bile acid conjugate: PD, pharmacology
- octanoic acid: CM, drug comparison
- octanoic acid: DV, drug development
- octanoic acid: PD, pharmacology
- decanoic acid: CM, drug comparison
- decanoic acid: DV, drug development
- decanoic acid: PD, pharmacology
- lauric acid: CM, drug comparison
- lauric acid: DV, drug development
- lauric acid: PD, pharmacology
- myristic acid: CM, drug comparison
- myristic acid: DV, drug development
- myristic acid: PD, pharmacology
- palmitic acid: CM, drug comparison
- palmitic acid: DV, drug development
- palmitic acid: PD, pharmacology
- stearic acid: CM, drug comparison
- stearic acid: DV, drug development
- stearic acid: PD, pharmacology
- oleic acid: CM, drug comparison

oleic acid: DV, drug development
 oleic acid: PD, pharmacology
 linoleic acid: CM, drug comparison
 linoleic acid: DV, drug development
 linoleic acid: PD, pharmacology
 arachidonic acid: CM, drug comparison
 arachidonic acid: DV, drug development
 arachidonic acid: PD, pharmacology
 icosapentaenoic acid: CM, drug comparison
 icosapentaenoic acid: DV, drug development
 icosapentaenoic acid: PD, pharmacology
 docosahexaenoic acid: CM, drug comparison
 docosahexaenoic acid: DV, drug development
 docosahexaenoic acid: PD, pharmacology
 amino acid: CM, drug comparison
 amino acid: DV, drug development
 amino acid: PD, pharmacology
 inorganic salt: CM, drug comparison
 inorganic salt: DV, drug development
 inorganic salt: PD, pharmacology
 human serum albumin
 omega 3 fatty acid: CM, drug comparison
 omega 3 fatty acid: DV, drug development
 omega 3 fatty acid: PD, pharmacology
 bendazac: CM, drug comparison
 bindarit: CM, drug comparison
 phenylbutazone: CM, drug comparison
 indometacin: CM, drug comparison
 antirheumatic agent: CM, drug comparison
 antirheumatic agent: DV, drug development
 antirheumatic agent: PD, pharmacology
 glycochenodeoxycholic acid: CM, drug comparison
 glycochenodeoxycholic acid: DV, drug development
 glycochenodeoxycholic acid: PD, pharmacology
 unindexed drug
 unclassified drug

RN (fish oil) 8016-13-5; (cholic acid) 32500-01-9, 361-09-1
 , 81-25-4; (deoxycholic acid) 83-44-3;
 (chenodeoxycholic acid) 474-25-9; (lithocholic acid)
 434-13-9; (octanoic acid) 124-07-2, 1984-06-1, 74-81-7; (decanoic
 acid) 334-48-5, 3398-75-2; (lauric acid) 115-05-9, 143-07-7; (myristic
 acid) 1715-79-3, 544-63-8; (palmitic acid) 57-10-3; (stearic acid)
 57-11-4, 646-29-7; (oleic acid) 112-80-1, 115-06-0; (linoleic acid)
 1509-85-9, 2197-37-7, 60-33-3, 822-17-3; (arachidonic acid) 506-32-1,
 6610-25-9, 7771-44-0; (icosapentaenoic acid) 25378-27-2, 32839-30-8;
 (docosahexaenoic acid) 25167-62-8, 32839-18-2; (amino acid) 65072-01-7;
 (human serum albumin) 9048-49-1; (bendazac) 20187-55-7; (phenylbutazone)
 129-18-0, 50-33-9, 8054-70-4; (indometacin) 53-86-1, 74252-25-8,
 7681-54-1; (glycochenodeoxycholic acid) 640-79-9
 CO Sigma (United States); Merck (Germany)

L51 ANSWER 4 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 96132330 EMBASE
 DN 1996132330
 TI Bile acid conjugation in early stage cholestatic liver disease
 before and during treatment with ursodeoxycholic acid.
 AU Fracchia M.; Setchell K.D.R.; Crosignani A.; Podda M.; O'Connell N.;
 Ferraris R.; Hofmann A.F.; Galatola G.
 CS Divisione di Gastroenterologia, Ospedale Mauriziano Umberto I, Largo
 Turati, 62, I-10128 Torino, Italy
 SO Clinica Chimica Acta, (1996) Vol. 248, No. 2, pp. 175-185.
 ISSN: 0009-8981 CODEN: CCATAR
 CY Netherlands
 DT Journal; Article
 FS 023 Nuclear Medicine

029 Clinical Biochemistry
 048 Gastroenterology
 037 Drug Literature Index

LA English
 SL English
 ED Entered STN: 960604
 Last Updated on STN: 960604

AB The efficiency of bile acid conjugation before and during therapy with 600 mg/day of ursodeoxycholic acid was measured in seven adult patients with early chronic cholestatic liver disease (6 with primary biliary cirrhosis; 1 with primary sclerosing cholangitis). Duodenal bile samples were obtained by aspiration and the proportion of unconjugated bile acids was determined using lipophilic anion exchange chromatography to separate bile acid classes, followed by analysis of individual bile acids by gas chromatography-mass spectrometry. The proportion of conjugated bile acids was determined by high-performance liquid chromatography. Use of a ^{99m}Tc -HIDA recovery marker permitted the absolute mass of unconjugated bile acids in the gallbladder to be calculated. Unconjugated bile acids comprised 0.4% of total biliary bile acids before and 0.2% during ursodeoxycholic acid therapy, indicating highly efficient conjugation of bile acids. During therapy, percentage unconjugated ursodeoxycholic acid significantly increased from (mean \pm S.D.) $13 \pm 13\%$ to $54 \pm 12\%$; $P < 0.002$. When the unconjugated and conjugated fractions of bile acids were compared, there was an enrichment in unconjugated fraction for cholic acid and ursodeoxycholic acid and a depletion for chenodeoxycholic acid both in basal condition and during ursodeoxycholic acid therapy, suggesting that hydrophilic bile acids were conjugated less efficiently. During therapy, the conjugation efficiency significantly increased for cholic acid and ursodeoxycholic acid. The pretreatment mass of total unconjugated bile acids in the gallbladder was (mean \pm S.D.) $4.4 \pm 3.2 \mu\text{mol}$, and was not significantly changed by ursodeoxycholic acid therapy ($6.2 \pm 3.5 \mu\text{mol}$). However, ursodeoxycholic acid therapy caused a significant increase in the mass of unconjugated ursodeoxycholic acid. It is concluded that endogenous bile acids and exogenous ursodeoxycholic acid when given at the usual dose are efficiently conjugated in patients with early cholestatic liver disease. Despite showing increased biliary unconjugated ursodeoxycholic acid during its oral administration, our data do not lend support to the occurrence of hypercholeresis due to cholehepatic shunting of bile acids.

CT Medical Descriptors:
 *cholestasis: DT, drug therapy
 *liver disease: DT, drug therapy
 article
 bile composition
 clinical article
 clinical trial
 gas chromatography
 high performance liquid chromatography
 human
 intravenous drug administration
 mass spectrometry
 oral drug administration
 priority journal
 Drug Descriptors:
 *bile acid conjugate: EC, endogenous compound
 *ursodeoxycholic acid: DT, drug therapy
 lidofenin tc 99m

RN (ursodeoxycholic acid) 128-13-2, 2898-95-5

L51 ANSWER 5 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 95240331 EMBASE
 DN 1995240331

TI Tauroursodeoxycholate increases rat liver ursodeoxycholate levels and limits lithocholate formation better than ursodeoxycholate.
 AU Rodrigues C.M.P.; Kren B.T.; Steer C.J.; Setchell K.D.R.
 CS Department of Pediatrics, Clinical Mass Spectrometry Center, Children's Hospital Medical Center, 3333 Burnet Avenue, Cincinnati, OH 45229, United States
 SO Gastroenterology, (1995) Vol. 109, No. 2, pp. 564-572.
 ISSN: 0016-5085 CODEN: GASTAB
 CY United States
 DT Journal; Article
 FS 037 Drug Literature Index
 048 Gastroenterology
 LA English
 SL English
 ED Entered STN: 950906
 Last Updated on STN: 950906
 AB Background and Aims: To explain the greater hepatoprotective effect of tauroursodeoxycholic acid vs. ursodeoxycholic acid, the absorption, hepatic enrichment, and biotransformation of these bile acids (250 mg/day) were compared in rats. Methods: Bile acids were determined in intestinal contents, feces, urine, plasma, and liver by gas chromatography-mass spectrometry. Results: The concentration of ursodeoxycholate in the liver of animals administered tauroursodeoxycholic acid (175 ± 29 nmol/g) was greater ($P < 0.05$) than in animals administered ursodeoxycholic acid (79 ± 19 nmol/g). Hepatic lithocholate was substantially higher after ursodeoxycholic acid administration (21 ± 10 nmol/g) than after tauroursodeoxycholic acid administration (12 ± 1 nmol/g). A concomitant reduction in the proportion of hydrophobic bile acids occurred that was greatest during tauroursodeoxycholic acid administration. In the intestinal tract, the mass of ursodeoxycholate and its specific metabolites was greater in rats administered tauroursodeoxycholic acid (27.2 mg) than those administered ursodeoxycholic acid (13.2 mg). In feces, the proportion of lithocholate was $21.9\% \pm 4.9\%$ and $5.4\% \pm 4.0\%$ after ursodeoxycholic acid and tauroursodeoxycholic acid administration, respectively. Conclusions: Compared with ursodeoxycholic acid, tauroursodeoxycholic acid induces a greater decrease in the percent composition of more hydrophobic bile acids within the pool, limits lithocholate formation, and increases hepatic ursodeoxycholate concentration. These differences are explained by increased hepatic extraction and reduced intestinal biotransformation and not by enhanced absorption of the amidated species.
 CT Medical Descriptors:
 *bile acid metabolism
 *biotransformation
 *liver protection
 amidation
 animal experiment
 animal tissue
 article
 blood level
 controlled study
 drug mechanism
 feces level
 gas chromatography
 hydrophobicity
 intestine absorption
 male
 mass spectrometry
 nonhuman
 oral drug administration
 priority journal
 rat
 urine level
 Drug Descriptors:
 *bile acid: EC, endogenous compound
 *lithocholic acid: EC, endogenous compound

*tauroursodeoxycholic acid: CM, drug comparison
 *tauroursodeoxycholic acid: PD, pharmacology
 *ursodeoxycholic acid: CM, drug comparison
 *ursodeoxycholic acid: PD, pharmacology
 bile acid conjugate: EC, endogenous compound
 RN (lithocholic acid) 434-13-9; (tauroursodeoxycholic acid)
 14605-22-2; (ursodeoxycholic acid) 128-13-2, 2898-95-5
 CO Sigma (United States)

L51 ANSWER 6 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 94299511 EMBASE
 DN 1994299511
 TI Effect of a medium dose of ursodeoxycholic acid with or without taurine
 supplementation on the nutritional status of patients with cystic
 fibrosis: A randomized, placebo-controlled, crossover trial.
 AU Merli M.; Bertasi S.; Servi R.; Diamanti S.; Martino F.; De Santis A.;
 Goffredo F.; Quattrucci S.; Antonelli M.; Angelico M.
 CS II Cattedra di Gastroenterologia, Viale dell'Universita, 37,00185 Rome,
 Italy
 SO Journal of Pediatric Gastroenterology and Nutrition, (1994) Vol. 19, No.
 2, pp. 198-203.
 ISSN: 0277-2116 CODEN: JPGND6
 CY United States
 DT Journal; Article
 FS 007 Pediatrics and Pediatric Surgery
 037 Drug Literature Index
 048 Gastroenterology
 LA English
 SL English
 ED Entered STN: 941027
 Last Updated on STN: 941027
 AB Ursodeoxycholic acid administration has been reported to improve
 cholestasis and inflammatory activity in primary biliary cirrhosis and, in
 an uncontrolled study, also in young adults with cystic fibrosis (CF) and
 chronic cholestasis. As an improvement in nutritional status was also
 observed in these young adult patients, we investigated whether the
 administration of a medium dose of ursodeoxycholic acid ameliorates the
 nutritional status of malnourished young adult CF patients with chronic
 liver disease. The study included 51 patients (27 male patients and 24
 female patients; age range, 8-32 years; median, 14) with body mass
 percentiles <90%. Patients were randomly assigned to receive either
 ursodeoxycholic acid (10- 12 mg/kg/day) alone or with taurine (18-22
 mg/kg/day). Patients were followed in a crossover fashion within each
 group; 6 months of treatment was randomly alternated with 6 months of
 placebo. Nine patients dropped out before concluding the study. Liver
 function tests, nutritional status, and coefficients of fat absorption
 were determined at entry and after each 6 months of placebo or treatment.
 Nutritional status and fat absorption were not significantly modified by
 either treatment. Liver function tests improved after ursodeoxycholic
 acid administration only in patients with concomitant chronic liver
 disease. Our findings indicate that 6 months of therapy with a medium
 dose of ursodeoxycholic acid, either alone or with taurine, does not
 improve the nutritional status of young malnourished CF patients. Higher
 doses given for longer periods might be worth investigating.

CT Medical Descriptors:
 *cholestasis: DT, drug therapy
 *cystic fibrosis: CN, congenital disorder
 *diet supplementation
 *nutritional status
 adolescent
 article
 body mass
 child
 clinical trial
 controlled study

crossover procedure
 dose response
 drug efficacy
 drug mixture
 enzyme therapy
 female
 human
 lipid absorption
 liver function test
 major clinical study
 male
 malnutrition: TH, therapy
 malnutrition: CO, complication
 oral drug administration
 pancreas insufficiency: DT, drug therapy
 pancreas insufficiency: CO, complication
 priority journal
 randomized controlled trial
 Drug Descriptors:
 *taurine: DT, drug therapy
 *taurine: CB, drug combination
 *taurine: CT, clinical trial
 *ursodeoxycholic acid: CT, clinical trial
 *ursodeoxycholic acid: DT, drug therapy
 *ursodeoxycholic acid: DO, drug dose
 *ursodeoxycholic acid: CB, drug combination
 alanine aminotransferase: EC, endogenous compound
 alkaline phosphatase: EC, endogenous compound
 amylase: DT, drug therapy
 amylase: CB, drug combination
 aspartate aminotransferase: EC, endogenous compound
 bile acid conjugate: EC, endogenous compound
 ceruletide
 ceruletide diethylamine
 feces lipid
 gamma glutamyltransferase: EC, endogenous compound
 pancrelipase: DT, drug therapy
 placebo
 tauroursodeoxycholic acid: EC, endogenous compound
 triacylglycerol lipase: CB, drug combination
 triacylglycerol lipase: DT, drug therapy
 trypsin: DT, drug therapy
 trypsin: CB, drug combination
 vitamin

RN (taurine) 107-35-7; (ursodeoxycholic acid) 128-13-2,
 2898-95-5; (alanine aminotransferase) 9000-86-6, 9014-30-6;
 (alkaline phosphatase) 9001-78-9; (amylase) 9000-90-2, 9000-92-4,
 9001-19-8; (aspartate aminotransferase) 9000-97-9; (ceruletide)
 17650-98-5; (ceruletide diethylamine) 71247-25-1; (gamma
 glutamyltransferase) 85876-02-4; (pancrelipase) 71060-52-1, 83869-36-7;
 (tauroursodeoxycholic acid) 14605-22-2; (triacylglycerol lipase)
 9001-62-1; (trypsin) 9002-07-7

CN (1) Takus

CO (1) Farmitalia carlo erba (Italy)

L51 ANSWER 7 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN

AN 92292458 EMBASE

DN 1992292458

TI Effect of ursodeoxycholic acid on the masses of biliary lipids and
 alkaline phosphatase within the gallbladder in chronic cholestatic liver
 disease.

AU Fracchia M.; Ferraris R.; Petrarulo M.; Secreto P.; Dunn T.; Galatola G.

CS Divisione di Gastroenterologia, Ospedale Mauriziano Umberto I, Largo
 Turati 62, I-10128 Torino, Italy

SO European Journal of Gastroenterology and Hepatology, (1992) Vol. 4, No.

10, pp. 843-848.
ISSN: 0954-691X CODEN: EJGHES
CY United Kingdom
DT Journal; Conference Article
FS 030 Pharmacology
037 Drug Literature Index
048 Gastroenterology
LA English
SL English
ED Entered STN: 921025
Last Updated on STN: 921025
AB Objectives: To verify whether the improvement of the cholestatic indices caused by ursodeoxycholic acid administered for chronic intrahepatic cholestasis is due to a dilution or a removal of the hydrophobic bile acids in the bile. To assess the effect of ursodeoxycholic acid on the masses in the gallbladder of other biliary lipids and alkaline phosphatase. Design: Open prospective study. Methods: Measurement of the masses of total bile acids, bile acid conjugates, cholesterol, phospholipid and alkaline phosphatase within the gallbladder in the fasting state before and after 4-6 weeks of therapy with 600 mg per day oral ursodeoxycholic acid in eight patients with chronic cholestatic liver disease. Results: Ursodeoxycholic acid caused a significant increase in the bile acid mass (from 1976 ± 593 to 4562 ± 1474 μmol ; $P < 0.02$), that was entirely due to an increased mass of its conjugates (from 35 ± 20 to 1623 ± 768 μmol ; $P < 0.05$), whereas the masses of all the other bile acid conjugates were not modified during therapy. In all eight patients, serum alkaline phosphatase concentration decreased during ursodeoxycholic acid therapy, whereas the alkaline phosphatase mass within the gallbladder increased, from 16 ± 3 IU to 35 ± 9 IU ($P < 0.02$). There was no change in the cholesterol and phospholipid masses. Conclusion: Our results indicate that the mechanism of action of ursodeoxycholic acid in chronic intrahepatic cholestasis is not mediated via a reduction of the hydrophobic bile acids handled by the liver, though these are diluted out by ursodeoxycholic acid. The finding of an increased mass of alkaline phosphatase in the gallbladder is probably due to the well known choleretic effect of ursodeoxycholic acid.
CT Medical Descriptors:
*cholestasis: DT, drug therapy
*chronic liver disease: DT, drug therapy
*gallbladder
*lipid bile level
adult
aged
alkaline phosphatase blood level
clinical article
conference paper
female
human
male
oral drug administration
primary biliary cirrhosis: DT, drug therapy
primary sclerosing cholangitis: DT, drug therapy
prospective study
Drug Descriptors:
*alkaline phosphatase: EC, endogenous compound
*ursodeoxycholic acid: DT, drug therapy
*ursodeoxycholic acid: PD, pharmacology
bile acid conjugate: EC, endogenous compound
cholesterol: EC, endogenous compound
phospholipid: EC, endogenous compound
RN (alkaline phosphatase) 9001-78-9; (ursodeoxycholic acid) 128-13-2
, 2898-95-5; (cholesterol) 57-88-5
CN Deursil
CO Labaz (Italy)

L51 ANSWER 8 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN

AN 90226900 EMBASE

DN 1990226900

TI Prevention of ursodeoxycholate hepatotoxicity in the rabbit by
conjugation with N-methyl amino acids.

AU Schmassmann A.; Hofmann A.F.; Angellotti M.A.; Ton-Nu H.-T.; Schteingart
C.D.; Clerici C.; Rossi S.S.; Rothschild M.A.; Cohen B.I.; Stenger R.J.;
Mosbach E.H.

CS Lipid Laboratory, Department of Surgery, Beth Israel Medical Center, 10
Nathan D. Perlman Place, New York, NY 10003, United States

SO Hepatology, (1990) Vol. 11, No. 6, pp. 989-996.
ISSN: 0270-9139 CODEN: HPTLD

CY United States

DT Journal; Article

FS 048 Gastroenterology
052 Toxicology
037 Drug Literature Index

LA English

SL English

ED Entered STN: 911213
Last Updated on STN: 911213

AB The effect of dietary administration of four different amino acid (N-acyl)
conjugates of ursodeoxycholic acid on biliary bile acid
composition, liver tests and hepatic morphology by light microscopy was
examined in the rabbit. Each group of four to five rabbits received a
chow diet supplemented with a single conjugate of
ursodeoxycholic acid ursodeoxycholyl-glycine, ursodeoxycholyl-sarcosine,
ursodeoxycholyl-taurine or ursodeoxycholyl-N-methyltaurine for 3 wks at a
dose of 50 mg/kg/day; a control group received chow alone. After 3 wks of
feeding, animals receiving ursodeoxycholyl-glycine or ursodeoxycholyl-
taurine had hepatotoxicity associated with abnormal liver tests.
Lithocholic acid made up $11\% \pm 2.7\%$ of biliary bile acids in the
ursodeoxycholyl-glycine and $10\% \pm 2.2\%$ in the ursodeoxycholyl-
taurine group. In contrast, animals receiving ursodeoxycholyl-sarcosine or
ursodeoxycholyl-N-methyltaurine had neither hepatotoxicity nor abnormal
liver tests and the proportion of lithocholic acid in biliary bile acids
increased much less. Complementary studies showed that
ursodeoxycholyl-sarcosine and ursodeoxycholyl-N-methyltaurine were not
biotransformed during hepatic transport and were resistant to
deconjugation and dehydroxylation in the rabbit. These
experiments indicate that the N-methyl amino acid conjugates of
ursodeoxycholic acid are nontoxic in the rabbit and resist
deconjugation and dehydroxylation. Such resistance decreases
formation of lithocholic acid in the colon, thus reducing its accumulation
and consequent induction of hepatotoxicity.

CT Medical Descriptors:
*conjugation
*liver toxicity: PC, prevention
drug conjugation
rabbit
animal experiment
nonhuman
male
oral drug administration
article
priority journal
Drug Descriptors:
*amino acid
*ursodeoxycholic acid: TO, drug toxicity

RN (amino acid) 65072-01-7; (ursodeoxycholic acid) 128-13-2,
2898-95-5

CO Diamalt aktiengesellschaft (Germany)

L51 ANSWER 9 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN

AN 89150062 EMBASE
 DN 1989150062
 TI The rapid evaluation of intestinal bacterial growth using a conjugate of ursodeoxycholic acid with para-aminobenzoic acid.
 AU Maeda Y.; Takahashi M.; Tashiro H.; Akazawa F.
 CS Department of Pharmacy, Chugoku Rosai Hospital, Hiroshima 737-01, Japan
 SO Journal of Pharmacobio-Dynamics, (1989) Vol. 12, No. 5, pp. 272-280.
 ISSN: 0386-846X CODEN: JOPHDQ
 CY Japan
 DT Journal
 FS 030 Pharmacology
 037 Drug Literature Index
 LA English
 ED Entered STN: 911212
 Last Updated on STN: 911212
 CT Medical Descriptors:
 *bacterial count
 *bacterial growth
 *blind loop syndrome
 *intestine flora
 animal model
 choloylglycine hydrolase
 rat
 microorganism
 animal experiment
 nonhuman
 male
 oral drug administration
 Drug Descriptors:
 bile acid
 4 ursodeoxycholamidobenzoic acid
 clindamycin
 glycocholic acid
 kanamycin
 paromomycin
 polymyxin b
 tinidazole
 vancomycin
 unclassified drug
 RN (clindamycin) 18323-44-9; (glycocholic acid) 475-31-0;
 (kanamycin) 11025-66-4, 61230-38-4, 8063-07-8; (paromomycin) 11035-13-5,
 1263-89-4, 1390-73-4, 51795-47-2, 54597-56-7, 7542-37-2, 84420-34-8;
 (polymyxin b) 1404-26-8, 1405-20-5; (tinidazole) 19387-91-8; (vancomycin)
 1404-90-6, 1404-93-9
 CO Wako pure chemical industry (Japan); Shionogi (Japan); Upjohn (Japan);
 Meiji seika kaisha (Japan); Kyowa hakko kogyo (Japan); Pfizer (Japan);
 Tokyo tanabe (Japan)
 L51 ANSWER 10 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 AN 88276787 EMBASE
 DN 1988276787
 TI HPLC assay of conjugated bile acids in gastric juice during ursodeoxycholic acid (Deursil®) therapy of bile reflux gastritis.
 AU Scalia S.; Pazzi P.; Stabellini G.; Guarneri M.
 CS Department of Pharmaceutical Sciences, University of Ferrara, 44100 Ferrara, Italy
 SO Journal of Pharmaceutical and Biomedical Analysis, (1988) Vol. 6, No. 6-8, pp. 911-917.
 ISSN: 0731-7085 CODEN: JPBADA
 CY United Kingdom
 DT Journal
 FS 029 Clinical Biochemistry
 048 Gastroenterology
 037 Drug Literature Index
 LA English

SL English
ED Entered STN: 911211
Last Updated on STN: 911211

AB A rapid high-performance liquid chromatographic method for the direct assay of the taurine and glycine conjugated bile acids in human gastric juice is described. After extraction with Sep-Pak C18 cartridges, compounds are baseline resolved on a reversed-phase column and detected by UV absorption. The procedure is linear from 10 μ mol l-1 to 1200 μ mol l-1, with recovery rates ranging from 87 to 100%. The present method is applicable to the quantification of bile acid conjugates in human gastric bile with satisfactory sensitivity, selectivity and precision. Intragastic bile acid compositions in 10 patients with bile reflux gastritis during Deursil® or placebo treatment are presented.

CT Medical Descriptors:
*bile reflux
*gastritis: DI, diagnosis
*gastritis: DT, drug therapy
*high performance liquid chromatography
*stomach juice
clinical article
human cell
human
methodology
oral drug administration
Drug Descriptors:
*bile acid conjugate
*ursodeoxycholic acid: DT, drug therapy

RN (ursodeoxycholic acid) 128-13-2, 2898-95-5
CN (1) Deursil
CO (1) Gipharmex (Italy)

L51 ANSWER 11 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN

AN 85080967 EMBASE
DN 1985080967
TI Synthesis, intestinal absorption and metabolism of sarcosine conjugated ursodeoxycholic acid.
AU Kimura M.; Hatono S.; Une M.; et al.
CS Institute of Pharmaceutical Sciences, Hiroshima University School of Medicine, Kasumi 1-2-3, Minami-Ku, Hiroshima 734, Japan
SO Steroids, (1984) Vol. 43, No. 6, pp. 677-687.
CODEN: STEDAM
CY United States
DT Journal
FS 037 Drug Literature Index
029 Clinical Biochemistry
023 Nuclear Medicine
048 Gastroenterology

LA English
ED Entered STN: 911210
Last Updated on STN: 911210

AB Sarcosine conjugated ursodeoxycholic acid (SUDC) was synthesized and its intestinal absorption and metabolism were studied in rat and hamster. Intestinal absorption study using bile fistula rat shows that more than 90% of SUDC administered intraduodenally was excreted in the bile within 24 hr. No change of the administered bile acid was seen during the absorption from the intestine, the passage of the liver, and the excretion into the bile. When [24-14C]SUDC and [11,12-3H2]-ursodeoxycholic acid were administered orally to a hamster, more than 95% of both the administered 14C and 3H were recovered from the feces within 6 days. Most (77%) of the fecal 14C-labeled compound was SUDC, whereas 95% of the fecal 3H-labeled compound was unconjugated lithocholic acid. These results indicate that SUDC, unlike taurine or glycine conjugated bile acid, resists bacterial deconjugation and 7-dehydroxylation.

CT Medical Descriptors:

*bile acid conjugation
 *drug absorption
 *drug bile level
 *drug distribution
 *drug elimination
 *drug feces level
 *drug identification
 *drug metabolism
 *drug monitoring
 *drug synthesis
 *drug tissue level
 *high performance liquid chromatography
 *infrared spectrometry
 *intestine absorption
 *ion exchange chromatography
 *nuclear magnetic resonance
 *sarcosodeoxycholic acid
 *sarcosodeoxycholic acid c 14
 *ursodeoxycholic acid c 14
 *ursodeoxycholic acid h 3
 metabolism
 priority journal
 drug analysis

oral drug administration

nonhuman

rat

small intestine

liver

animal experiment

Drug Descriptors:

*lithocholic acid

*sarcosine

radioisotope

RN (lithocholic acid) 434-13-9; (sarcosine) 107-97-1

CO Daiichi; Nen

L51 ANSWER 12 OF 12 EMBASE COPYRIGHT 2005 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN

AN 84147710 EMBASE

DN 1984147710

TI Effect of ursodeoxycholate and its taurine conjugate on bile
acid synthesis and cholesterol absorption.

AU Hardison W.G.M.; Grundy S.M.

CS Department of Medicine, Veterans Administration Medical Center, San Diego,
TX, United States

SO Gastroenterology, (1984) Vol. 87, No. 1, pp. 130-135.

CODEN: GASTAB

CY United States

DT Journal

FS 037 Drug Literature Index

048 Gastroenterology

006 Internal Medicine

029 Clinical Biochemistry

003 Endocrinology

023 Nuclear Medicine

LA English

ED Entered STN: 911210

Last Updated on STN: 911210

AB Six male subjects with normal gastroenterologic function were studied to
determine the effects of ursodeoxycholic (15 mg/kg·day) and
tauroursodeoxycholic (20 mg/kg·day) acid feeding on bile acid
synthesis and cholesterol absorption. Each bile acid was fed for 1 mo and
withheld for the next month. Subjects remained on a metabolic ward and
consumed a constant diet of 500 mg of cholesterol mixed with solid and
liquid formulas. Before the study started, each subject received 50
µCi of [4-14C]cholesterol intravenously. During the study, stools were

collected for the determination of 24-h fecal acidic and neutral sterol excretion, and blood was drawn twice weekly for determination of serum cholesterol specific activity. At the end of each month an intestinal perfusion study was performed to measure total bile acid pool size and hourly biliary secretion rates of cholesterol, phospholipid, and bile acid. From these data, the percentage of cholesterol absorption and total endogenous bile acid synthesis could be calculated. Neither ursodeoxycholic nor tauroursodeoxycholic acid feeding decreased endogenous bile acid synthesis. During bile acid feeding periods, the percentage of cholesterol absorption was decreased.

CT Medical Descriptors:

*cholesterol c 14
 *drug efficacy
 *intestine absorption
 oral drug administration

human

normal human

liver

human experiment

Drug Descriptors:

*bile acid
 *cholesterol
 *tauroursodeoxycholic acid
 *ursodeoxycholic acid
 radioisotope

RN (cholesterol) 57-88-5; (tauroursodeoxycholic acid) 14605-22-2;

(ursodeoxycholic acid) 128-13-2, 2898-95-5

CO Nen (United States)

=> b biosis

FILE 'BIOSIS' ENTERED AT 09:39:27 ON 24 JUN 2005

Copyright (c) 2005 The Thomson Corporation

FILE COVERS 1969 TO DATE.

CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNs) PRESENT
 FROM JANUARY 1969 TO DATE.

RECORDS LAST ADDED: 23 June 2005 (20050623/ED)

FILE RELOADED: 19 October 2003.

=> d all 160 tot

L60 ANSWER 1 OF 1 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN

AN 2003:31809 BIOSIS

DN PREV200300031809

TI Absorption of biologically active peptide hormones from the small intestine of rat.

AU Wheeler, S.; McGinn, B. J.; Lucas, M. L.;

Morrison, J. D. [Reprint Author]

CS University of Glasgow, West Medical Building, Glasgow, G12 8QQ, UK

SO Acta Physiologica Scandinavica, (November 2002) Vol. 176, No. 3, pp. 203-213. print.

ISSN: 0001-6772 (ISSN print).

DT Article

LA English

ED Entered STN: 8 Jan 2003

Last Updated on STN: 8 Jan 2003

AB Absorption of the 4, 10 and 34 amino acid forms of gastrin from the small intestine has been investigated in anaesthetized rats. The method of assessment of successful absorption of the hormone into the systemic circulation was when the amount of acid secreted by the stomach over consecutive 15-min periods was increased. When the natural hormones were infused into the ileum in a relatively high dose, there was no increase in gastric acid secretion, indicating that they had not been absorbed. Each

of the forms of gastrin was conjugated at the free amino terminus to the carboxyl group of cholic acid. Subsequent infusion of the conjugated form of gastrin into the ileum, this time in relatively low doses, resulted in substantial and prolonged increases in gastric acid secretion, indicating that these hormones had been successfully absorbed. In addition, conjugation of the 10 and 34 amino acid forms of gastrin with cholic acid was shown to increase markedly the potency in evoking an increase in gastric acid secretion in response to intravenous injection of the hormone. Absorption of the gastrin conjugates was specific to the ileum thus indicating that they had been absorbed through the bile salt transporters.

CC Biochemistry studies - General 10060
 Digestive system - Physiology and biochemistry 14004
 IT Major Concepts
 Biochemistry and Molecular Biophysics; Digestive System (Ingestion and Assimilation)
 IT Parts, Structures, & Systems of Organisms
 ileum: digestive system; small intestine: digestive system; stomach: digestive system
 IT Chemicals & Biochemicals
 bile salt transporters; biologically active peptide hormones: absorption; cholic acid; gastrin conjugates: absorption
 ORGN Classifier
 Muridae 86375
 Super Taxa
 Rodentia; Mammalia; Vertebrata; Chordata; Animalia
 Organism Name
 Wistar rat (common): male
 Taxa Notes
 Animals, Chordates, Mammals, Nonhuman Vertebrates, Nonhuman Mammals, Rodents, Vertebrates
 RN 81-25-4 (cholic acid)

=> b home

FILE 'HOME' ENTERED AT 09:39:33 ON 24 JUN 2005

=>